

Exercise Recommendations for Individuals with Spinal Cord Injury

Patrick L. Jacobs^{1,2} and Mark S. Nash^{1,3}

- 1 Department of Neurological Surgery, University of Miami School of Medicine, Miami, Florida, USA
- 2 Center of Excellence for Restoration of Function in Chronic SCI, Miami VA Medical Center, Miami, Florida, USA
- 3 Department of Orthopedics & Rehabilitation, University of Miami School of Medicine, Miami, Florida, USA

Contents

Abstract	728
1. Neurological Classification of Persons with Spinal Cord Injury (SCI)	729
1.1 Upper and Lower Motor Neuron Lesions	730
1.2 Defining Spinal Cord Lesions: Plegias	731
1.3 Neurological, Sensory and Motor Levels	731
1.4 Complete and Incomplete Spinal Cord Lesions	731
1.5 SCI Syndromes	732
1.5.1 Anterior Cord Syndrome	732
1.5.2 Central Cord Syndrome	732
1.5.3 Brown-Sequard Syndrome	733
1.5.4 Conus Medullaris and Cauda Equina Syndrome	733
2. Medical and Health Consequences of SCI	733
2.1 Physical Deconditioning	733
2.2 Musculoskeletal Decline	734
2.3 Cardiovascular Disease, Dyslipidaemia and Comorbidities	734
2.4 Cardiac Structure and Function	735
2.5 Vascular Structure and Function	735
2.6 Autonomic Dysregulation	736
3. Volitional Upper Extremity Exercise in Persons with SCI	736
3.1 Acute Physiological Responses to Exercise Stress	736
3.2 Cardiorespiratory Testing	737
3.3 Endurance Training	738
3.4 Resistance Training for Persons with SCI	739
4. Electrically Stimulated Exercise in Persons with SCI	740
4.1 Basis of Electrically Stimulated Movement	740
4.2 Leg Cycling Exercise	740
4.3 Bipedal Ambulation	741
5. Risks of Exercise When Performed by Persons with SCI	742
5.1 Autonomic Dysreflexia	742
5.2 Musculoskeletal Injury	743
5.3 Hypotension	743
5.4 Thermal Dysregulation	743
6. Conclusions	743

Abstract

Persons with spinal cord injury (SCI) exhibit deficits in volitional motor control and sensation that limit not only the performance of daily tasks but also the overall activity level of these persons. This population has been characterised as extremely sedentary with an increased incidence of secondary complications including diabetes mellitus, hypertension and atherogenic lipid profiles. As the daily lifestyle of the average person with SCI is without adequate stress for conditioning purposes, structured exercise activities must be added to the regular schedule if the individual is to reduce the likelihood of secondary complications and/or to enhance their physical capacity. The acute exercise responses and the capacity for exercise conditioning are directly related to the level and completeness of the spinal lesion. Appropriate exercise testing and training of persons with SCI should be based on the individual's exercise capacity as determined by accurate assessment of the spinal lesion. The standard means of classification of SCI is by application of the *International Standards for Classification of Spinal Cord Injury*, written by the Neurological Standards Committee of the American Spinal Injury Association. Individuals with complete spinal injuries at or above the fourth thoracic level generally exhibit dramatically diminished cardiac acceleration with maximal heart rates less than 130 beats/min. The work capacity of these persons will be limited by reductions in cardiac output and circulation to the exercising musculature.

Persons with complete spinal lesions below the T₁₀ level will generally display injuries to the lower motor neurons within the lower extremities and, therefore, will not retain the capacity for neuromuscular activation by means of electrical stimulation. Persons with paraplegia also exhibit reduced exercise capacity and increased heart rate responses (compared with the non-disabled), which have been associated with circulatory limitations within the paralysed tissues. The recommendations for endurance and strength training in persons with SCI do not vary dramatically from the advice offered to the general population. Systems of functional electrical stimulation activate muscular contractions within the paralysed muscles of some persons with SCI. Coordinated patterns of stimulation allows purposeful exercise movements including recumbent cycling, rowing and upright ambulation. Exercise activity in persons with SCI is not without risks, with increased risks related to systemic dysfunction following the spinal injury. These individuals may exhibit an autonomic dysreflexia, significantly reduced bone density below the spinal lesion, joint contractures and/or thermal dysregulation. Persons with SCI can benefit greatly by participation in exercise activities, but those benefits can be enhanced and the relative risks may be reduced with accurate classification of the spinal injury.

The human spinal cord is a complex association of upper and lower motor neurons that functions as a bidirectional conduit between the brain and its motor, sensory and autonomic targets. It also serves as a site for reflex integration between body sensors and their motor and autonomic effectors. Because spinal cord functions differ by level and structure, injury to

or disease of spinal tracts results in varying types and degrees of dysfunction depending upon the specific neural structures affected.

The interruption of spinal cord functions by trauma affects 10 000 Americans annually, with an estimated 179 000 persons having survived their initial injury.^[1-4] Thereafter, these individuals experience

unique physical, social and psychological changes throughout their lives, including diminished ability to perform and benefit from exercise conditioning.^[5,6] The latter limitation is cause for concern as: (i) individuals with spinal cord injury (SCI) are usually young and physically active at the time of injury;^[3] (ii) profound physical deconditioning is common after injury;^[7-9] and (iii) physical deconditioning contributes to multisystem medical complications,^[10-19] activity limitations^[7,20,21] and accelerated aging.^[6,7,22-26]

Many reviews and chapters published over the past decades have addressed the need for persons with SCI to adopt habitual exercise as part of a healthy lifestyle.^[5,6,9,17,18,27-33] Exercise options contained in these reports have focused on use of muscle activities still under voluntary nervous system control, as well as sequenced contractions of muscles stimulated by electrical current. Outcomes of studies cited in these monographs provide credible evidence that exercise performed by persons with SCI enhances physical conditioning and reduces multisystem disease susceptibility. They further suggest that habitual exercise might reduce fatigue, pain, weakness, joint deterioration and incipient neurological deficits that appear as persons age with disability. As these deficits challenge the ability of those with SCI to perform essential daily activities first mastered after injury, their prevention will likely foster fullest health and life satisfaction for those aging with a disability.

Use of exercise as a healthy activity and lifestyle elective for those with SCI might appear fairly straightforward. To the contrary, exercise options available to those with SCI are more limited than those without disability,^[33] their acute exercise and training responses less robust than those of persons without SCI,^[28,34,35] and the risks of impudent activity greater and longer lasting.^[33,36] This makes an understanding of neurological classification, exercise opportunities and risks of activity important if those with SCI are to benefit and not be harmed from exercise conditioning. This paper will review common medical problems experienced by persons with SCI that lend themselves to benefit from exer-

cise conditioning, contemporary methods of injury classification, exercise options and established benefits of training, and risks imposed by inappropriate exercise recommendations for those with SCI.

1. Neurological Classification of Persons with Spinal Cord Injury (SCI)

If trauma to, or disease of, the spinal cord always resulted in its anatomical or physiological transection, classification of spinal cord dysfunctions following SCI would be relatively easy. However, the spinal cord is rarely severed unless penetrated by a bullet or severed by its bony covering during very high velocity impact. More often the cord remains anatomically intact but suffers contusion, infarction or mechanical deformation that interrupts its local or relay circuitry. It also can undergo secondary damage from inflammatory autodestruction, which explains the routine administration of high-dose corticosteroids within hours of an SCI.^[37] Notwithstanding the cause of spinal damage, more than half of the survivors will experience varying degrees of motor, sensory or autonomic sparing at different spinal cord levels,^[2,38,39] making classification of persons with SCI a challenging, yet important, science.^[4]

To assist with accurate and consistent classification, uniform standards have been developed that allow persons with SCI to be systematically examined and classified, and to document changes in sensorimotor function that accompany the passage of time, clinical treatments or research interventions. The benchmark system used to classify persons with SCI is the *International Standards for Classification of Spinal Cord Injury* (revised 2000), which is written by the Neurological Standards Committee of the American Spinal Injury Association (ASIA), and endorsed as the recommended international standards by the International Spinal Cord Society (ISCoS; formerly the International Medical Society for Paraplegia, and hereinafter the 'ASIA Guidelines') [figure 1].^[40] While a complete description of spinal column and spinal cord anatomy is beyond the scope of this review, the accurate description of an injury to a given spinal cord segment involves designation of a spinal column region




Fig. 1. American Spinal Injury Association (ASIA) classification of spinal cord injury (reproduced from the American Spinal Injury Association,^[41] with permission).

(e.g. cervical, thoracic, lumbar or sacral), a spinal nerve within that region, and the degree to which the injury is neurologically complete or incomplete. Should the injury result in a specific spinal cord syndrome this should also be designated.

1.1 Upper and Lower Motor Neuron Lesions

It is clinically important for those working with subjects having SCI to differentiate between upper and lower motor neuron injuries. The spinal cord is organised in such a way that either upper or lower motor neuron function, or both, can be disrupted by

trauma or disease. The long tracts of the spinal cord are upper motor neurons. By convention, the lower motor neuron system begins in the anterior horn of the grey matter and exits the intervertebral foramen to join the returning sensory axons as a mixed nerve. The motor portion of this mixed nerve then travels to its target muscle(s). The returning sensory nerve fibres travel in this mixed nerve but, upon entry to the foramen, form a dorsal root that penetrates the cord in the dorsal root entry zone and synapses with the anterior horn cells of the anterior grey matter. This completes the motor-sensory reflex arc. Lesions of upper motor neurons result in ‘decentralisa-

tion' of the nervous system, spastic paralysis and exaggerated sensorimotor reflexes below the injury. This means that motor, sensory and autonomic reflex activities are preserved, but no longer under command of the brain. The contrasting injury is a lower motor neuron lesion, which often accompanies SCI at the T₁₀ level or lower and almost always occurs at, or caudal to, T₁₂. Persons with these lesions lose central nervous system control of sensorimotor functions, as well as sensorimotor reflex activity, which causes flaccid paralysis and areflexia ('denervation'). This explains the greater loss of lower extremity muscle mass in individuals with flaccid rather than spastic paralysis. Otherwise, injury to the cord at any site resulting in damage to the reflex arc can leave a denervated (areflexic) segment among neighbouring segments that remain spastic.

It is important to note that while spastic muscles are capable of contracting upon stimulation with transcutaneous electrical alternating currents, generally, flaccid muscles are not. This provides a means of discriminating upper from lower motor neuron lesions and also determining subject or patient candidacy for use of electrical prostheses that provide exercise or ambulation in persons with SCI.

1.2 Defining Spinal Cord Lesions: Plegias

Individuals with SCI are described as having lesions affecting either sensorimotor function of all limbs or the upper extremities alone. The former are said to have tetraplegia (preferred to 'quadriplegia'), which is defined as: "A term referring to impairment or loss of motor and/or sensory function in the cervical segments of the spinal cord due to damage of neural elements within the spinal canal. Tetraplegia results in impairment of function in the arms as well as in the trunk, legs and pelvic organs. It does not include brachial plexus lesions or injury to peripheral nerves outside the neural canal."^[42,43] Paraplegia is defined as: "A term referring to impairment or loss of motor and/or sensory function in the thoracic, lumbar or sacral (but not cervical) segments of the spinal cord, secondary to damage of neural elements within the spinal canal. With paraplegia, arm functioning is spared, but, depending on

the level of injury, the trunk, legs and pelvic organs may be involved. The term is used when referring to cauda equina and conus medullaris injuries, but not to lumbosacral plexus lesions or injury to peripheral nerves outside the neural canal."^[42,43]

1.3 Neurological, Sensory and Motor Levels

Neurological level refers to the most caudal segment of the spinal cord with normal sensory and motor function on both sides of the body. Because the results of sensorimotor testing often differ by side, up to four different segments may be identified in determining the neurological level, i.e. R-sensory, L-sensory, R-motor and L-motor. In such cases it is strongly recommended that each of these segments be separately recorded and that a single 'level' not be used, as this can be misleading. When the term 'sensory level' is used, it refers to the most caudal segment of the spinal cord with normal sensory function on both sides of the body; the 'motor level' is similarly defined with respect to motor function. These 'levels' are determined by neurological examination of: (i) a key sensory point within each of 28 dermatomes on the right and 28 dermatomes on the left side of the body;^[42] and (ii) a key muscle within each of ten myotomes on the right and ten myotomes on the left side of the body. Assessment of sensory and motor level is key in those with paraplegia having injuries from T₁ through T₈, whereas motor function is the determining factor in classification of tetraplegia.

1.4 Complete and Incomplete Spinal Cord Lesions

Confusion and imprecise language surround the definitions of neurological completeness and incompleteness. Historically, the Functional Classification scale of Frankel et al.^[44] was used to assign a grade of SCI from A to E. These grades were assigned on the basis of sensorimotor sparing below the level of SCI. In contrast, the ASIA scale uses 'sacral sparing' as the criterion for determining neurological completeness.^[39,42,45] This requires a test of motor function examining the presence of voluntary contraction of the external anal sphincter upon digital

examination. The ASIA definitions of complete and incomplete lesions are:

- ‘Complete injury’: a term describing absence of sensory and motor function in the lowest sacral segment.^[42]
- ‘Incomplete injury’: a term describing partial preservation of sensory and/or motor functions below the neurological level *and* including the lowest sacral segment. Sacral sensation includes sensation at the anal mucocutaneous junction as well as deep anal sensation.^[42]

Both ASIA and Frankel methods are useful for describing completeness of injuries, as an individual with infralesional sensory sparing (i.e. incomplete sensory loss; Frankel Grade B) can still be neurologically complete when using the ASIA guidelines. Also, the ASIA C classification is so broad as to sometimes limit useful classification of motor and functional abilities of patients with SCI that may be better understood when using Frankel grades.

Accurate descriptions of complete and incomplete lesions are often confounded by recovery of sensorimotor function within weeks or months after injury, which is not uncommon for some individuals initially presenting without preserved reflex or sensorimotor functions.^[46] This recovery, however, will more likely occur in individuals with very low velocity injuries in which cord bleeding or anatomic disruption of neural elements are limited. Several mechanisms have been postulated to explain ‘return of function’ after SCI. The first involves a belief that a ‘spinal shock’ occurs immediately upon SCI in some individuals – a state of areflexia thought to result from hyperpolarisation of cord neurons accompanying injury-related loss of descending facilitation.^[47] Thereafter, those with incomplete lesions often recover reflexes in a caudal to rostral sequence, although the extent of reflex recovery has questionable value in predicting eventual patterns or extent of motor recovery.^[47] Otherwise, post-synaptic receptor up-regulation and synaptic growth have been postulated as explanations for neuronal recovery following SCI,^[48,49] although the precise mechanisms promoting these changes remain incompletely understood. Notwithstanding the controversial ex-

planations for resolution of ‘spinal shock’ and recovery of sensorimotor functions, the designation of ‘neurological completeness’ within the first year of injury leaves open the possibility for erroneous classification of a subject thereafter. This explains why many investigators use the first anniversary of injury as both a benchmark for neurological stability and a criterion for enrolment of subjects with SCI in research studies, especially when sensorimotor function is under study. Clearly, however, the use of the term ‘complete’ written on an initial trauma examination might ultimately have little bearing on whether recovery of reflexes or sensorimotor function might result. Thus, examination and classification at the beginning of exercise testing and training is needed.

1.5 SCI Syndromes

As the various spinal columns transmit different types of information, and because most persons surviving SCI do not experience complete severance of their cords, many different patterns of spared sensorimotor function are possible. The syndromes that ensue are usually described when defining the levels and completeness of injury. The major syndromes, their definitions and causations are listed in the following sections.

1.5.1 Anterior Cord Syndrome

An anterior cord syndrome is “a lesion that produces variable loss of motor function and of sensitivity to pain and temperature, while preserving proprioception”.^[42] An anterior cord syndrome results primarily in a profound motor loss below the level of injury. Destruction of the anterior portions of the white and grey matter of the spinal cord affects the corticospinal motor tracts and, to a lesser extent, sensory tracts that mediate light touch and pressure.

1.5.2 Central Cord Syndrome

A central cord syndrome is “a lesion occurring almost exclusively in the cervical region, that produces sacral sensory sparing and greater weakness in the upper limbs than in the lower limbs”.^[42] A person with a central cord syndrome often presents

with greater motor impairment of the upper than lower extremities if the damage or disease is primarily limited to the central grey and white spinal cord matter. Because upper extremity motor function is primarily activated by the medial corticospinal tracts, sensorimotor functions of the lower extremities are usually affected to a lesser degree. Central cord injury is common among individuals having congenitally narrow spinal canals and in those, especially aging persons, whose osteoarthritic pathology results in spinal canal stenosis.

1.5.3 Brown-Sequard Syndrome

A Brown-Sequard syndrome is “a lesion that produces relatively greater ipsilateral proprioceptive and motor loss and contralateral loss of sensitivity to pain and temperature”.^[42] The Brown-Sequard syndrome results from an anterior-posterior hemisection of the spinal cord often accompanying a penetrating wound that severs the neural elements. When the elements of only one lateral portion of the spinal cord are destroyed, the injury results in an unusual pattern of sensorimotor function with a loss of infralesional motor function, proprioception, fine touch and vibration discrimination on the same side of injury, but loss of pain, temperature, crude touch and deep pressure on the opposite side of injury. The syndrome is explained by decussation of motor tracts in the brain stem before they descend in the cord. In contrast, ascending sensory tracts of the anterolateral systems cross to the opposite side at (or near) the level in which the dorsal roots enter the cord. This explains why ‘same side’ motor deficits are common in patients with Brown-Sequard syndrome, but sensory preservation is greater on the contralateral side of injury.

1.5.4 Conus Medullaris and Cauda Equina Syndrome

A conus medullaris syndrome involves an “injury of the sacral cord (conus) and lumbar nerve roots within the spinal canal, which usually results in an areflexic bladder, bowel and lower limbs”.^[42] Sacral segments may occasionally show preserved reflexes (e.g. bulbocavernosus and micturition reflexes). A cauda equina syndrome involves “injury to the lum-

bosacral nerve roots within the neural canal resulting in areflexic bladder, bowel and lower limbs”.^[42]

2. Medical and Health Consequences of SCI

Persons with SCI face unique health challenges throughout their lives, and their injuries dissociate the normally well integrated homeostatic responses of body systems known to accompany physical activity. Nervous system damage disrupts to varying degrees the necessary signal integration among motor, sensory and autonomic targets, and thus has a profound effect on fitness, exercise capacities and health. Depending on the level and type of cord lesion, persons with SCI are among the most physically deconditioned of all humans.^[7,50] Not surprisingly, young persons with chronic SCI experience accelerated pathological states and conditions normally associated with physical deconditioning and premature aging, including: dyslipidaemias and heart disease;^[23,24,51,52] arterial circulatory insufficiency^[53-57] and clotting disorders;^[58,59] bone and joint diseases;^[60-62] and pain of musculoskeletal and neuropathic origins.^[63-68]

Notwithstanding their physical limitations, many persons with SCI can still undergo exercise reconditioning. Those who retain upper extremity function can participate in a wide variety of exercise activities and sports,^[6,69] and ambulate with the assistance of orthoses and computer-controlled electrical neuroprostheses.^[70-76] Individuals with upper motor neuron lesions can pedal cycle ergometers by surface electrical stimulation of selected lower extremity muscle groups under computer control.^[30,31,33] Many body organs and tissues acutely respond to exercise despite their decentralised or denervated states, and because many survivors of SCI experience complete sensory loss or significantly diminished nociceptive responses, electrically stimulated muscle contractions can often be utilised without pain.

2.1 Physical Deconditioning

Persons with SCI usually live sedentary lives,^[7,9,68,77] which explains their poor physical fit-

ness and heightened risk of cardiovascular morbidity and mortality.^[9,22,51,78] Nearly one in four healthy young persons with paraplegia fails to achieve levels of oxygen consumption ($\dot{V}O_2$) on an arm exercise test sufficient to perform many essential activities of daily living.^[8] While those with paraplegia have far greater capacities for activity and more extensive choices for exercise participation than persons with tetraplegia, they are only marginally more fit.^[7,11]

2.2 Musculoskeletal Decline

Altered structural and contractile properties of muscle after SCI limit the ability of totally paralysed and weakened muscle to sustained intense contractions for extended durations. Most studies of sublesional muscle after SCI in humans report fibres that:

- are smaller than those above the lesion and those of persons without SCI;^[79-83]
- have less contractile protein;^[84]
- produce lower peak contractile forces;^[85,86]
- transform toward fast phenotypic protein expression;^[81,87-90]
- increase myosin heavy chain isoforms;^[90,91]
- decrease their resistance to fatigue.^[19,79,84,92,93]

Muscle fibre cross-sectional area declines within 1 month of SCI,^[80,89] while electrical stimulation of muscle paralysed for more than 1 year evokes forces only one-seventh to one-third those of subjects without SCI.^[85,86,94] Physiological and contractile properties adopted after SCI further compromise the ability of muscle without motor sparing to optimally increase in size and metabolic activity after exercise reconditioning,^[95] which has adverse consequences for cardiovascular, musculoskeletal and comorbid diseases to which those with SCI are highly susceptible.

Persons sustaining SCI develop muscles below the level of the lesion that are either hypertonic or atonic depending on the level and type of SCI. Hypertonia is the more common condition in which exaggerated rate-dependent stretch results in spastic contraction. This results from damage to the upper motor neurons alone, in which descending inhibition of reflex response is interrupted by cord injury. Hypertonia in these individuals can be worsened by

a number of 'noxious' stimuli unrelated to muscle stretch, including urinary voiding, venous thrombosis, thermal dysregulation, occult fracture or infection.^[96] In contrast, damage to lower motor neurons involving injury below T₁₀ usually results in flaccid paralysis and loss of neuromuscular response to administration of alternating electrical currents.

Rapid bone demineralisation is expected during the first year after SCI, after which bone density levels continue to slowly decay. Increased urinary excretion of calcium and hydroxyproline,^[97] and progressive rarefying of bone on radiographs are evident throughout this period. About one-third to one-half of bone mineral density is lost by 1 year after injury, with primary losses occurring in the supracondylar femur. The inevitable course of SCI leads to underhydroxylated and hypocalcific bone^[98,99] with permanently heightened susceptibility to fracture, even following trivial or imperceptible trauma.^[100] Joints experience similar deterioration^[60] and heightened injury susceptibility brought on by cartilage atrophy^[101] and joint space deformities.^[102-104]

2.3 Cardiovascular Disease, Dyslipidaemia and Comorbidities

Epidemiological studies conducted in the early 1980s, and thereafter, reported emergence of cardiovascular disease as a major cause of death in persons with SCI.^[1,3,24,105] While genitourinary complications accounted for 43% of deaths in the 1940s and 1950s, mortality from these causes were reduced to 10% of cases in the 1980s and 1990s.^[24,106,107] Cardiovascular diseases currently represent the most frequent cause of death among persons surviving >30 years after injury (46% of deaths) and among persons >60 years of age (35% of deaths)^[107,108]

Decline of cardiovascular function in persons aging with SCI mirrors that experienced by persons who age without SCI, although at an accelerated rate.^[23] Asymptomatic cardiovascular disease also occurs at earlier ages after SCI^[108,109] and its symptoms may be masked by interruption of ascending afferent pain fibres conveying warnings of impending heart damage or death.^[108-110] Several major risk

factors commonly reported in persons with SCI have been linked with their accelerated course of cardiovascular disease; these include dyslipidaemia^[7,22,78] and a sedentary lifestyle imposed by muscle paralysis and limited exercise options.^[8,9] The cardiovascular disease risks of individuals without SCI are worsened by hyperinsulinaemia^[22,111,112] and elevated percentages of body fat,^[113,114] which are common among persons with SCI.^[24]

An atherogenic lipid profile is commonly reported in persons with chronic SCI.^[22,78,112-115] This lipid profile satisfies criteria for designation as 'dyslipidaemic', i.e. elevated total cholesterol (TC), triglyceridaemia and plasma low density lipoprotein-cholesterol (LDL-C) or depressed high density lipoprotein-cholesterol (HDL-C).^[116] All have all been reported in sedentary persons with SCI, although elevated TC is not observed in all of these studies,^[115,117] and LDL-C concentrations, while sometimes elevated, show patterns of elevation typical of those reported in the general population.^[115] The most consistent finding of reports examining the lipid profiles of persons with SCI is a depressed blood plasma concentration of the cardioprotective HDL-C^[22,78,114,115,118,119] whose levels are inversely associated with cardiovascular risk.^[120] More than 40% of persons with SCI have HDL-C levels below the earlier criterion score for high cardiovascular risk (HDL >35 mg/dL). When combined with other known risks factors for vascular disease in persons with SCI (e.g. prevalent truncal obesity,^[114] elevated body mass indices,^[113] physical inactivity^[8,9] reduced lean body mass,^[95,113,121-123] diabetes,^[124,125] metabolic syndrome X^[126] and advancing age),^[18] the risks of abnormal lipid profile on disease progression become magnified.

Insulin resistance occurring in a high percentage of persons with SCI was first reported more than two decades ago.^[125] Since that time, others have confirmed this finding and included insulin resistance among the cardiovascular risks sustained by persons aging with SCI.^[22,111,112,114,127] Almost half the persons with SCI live in a state of carbohydrate intolerance or insulin resistance.^[22,125] A reason for prevalent insulin resistance in persons with SCI have not

been firmly identified, although physical inactivity,^[127] truncal obesity^[113,114] and sympathetic dysfunction^[111,112] have been suggested as causes. An association may also exist between their abnormal lipid profiles and insulin resistance. Such an association has been identified in persons without SCI having depressed serum HDL-C, as they are also especially prone to insulin resistance.^[128-130] This risk profile closely matches that of persons with SCI, in whom isolated low HDL-C and insulin resistance are often comorbid.^[51]

2.4 Cardiac Structure and Function

Individuals with chronic SCI experience various types of circulatory dysregulation depending on the level of their cord lesion.^[96] When injury occurs above the level of sympathetic outflow at the T₁ spinal level, resting hypotension with mean arterial pressures of 70mm Hg are common.^[131] In addition to challenging effective orthostatic pressure regulation, low pressures also instigate altered heart structure and function observed after SCI.^[132] As size and architecture of the human heart are known to be influenced by peripheral circulatory volume and systemic pressures, withdrawal from normal activity levels and altered circulatory dynamics transform the structure of the heart and alter its pumping efficiency.^[133] For those with tetraplegia, a chronic reduction of cardiac preload and myocardial volume coupled with pressure underloading causes the left ventricle to atrophy.^[132,134] In contrast, long-term survivors of paraplegia are normotensive and have normal left ventricular mass and resting cardiac output, but experience a cardiac output comprised of elevated resting heart rate (HR) and depressed resting stroke volume (SV).^[28,134] This lowered SV is attributed to decreased venous return from the immobile lower extremities or frank venous insufficiency of the paralysed limbs.^[135,136]

2.5 Vascular Structure and Function

The volume and velocity of lower extremity arterial circulation is significantly diminished after SCI, with volume flow of about half to two-thirds that reported in matched subjects without paraly-

sis.^[56,137] This circulatory 'hypokinesia'^[138] results from loss of autonomic control of blood flow as well as diminished regulation of local blood flow by vascular endothelium.^[56] The lowering of volume and velocity contribute to heightened thrombosis susceptibility commonly reported in those with acute and subacute SCI.^[58] A contributing factor to thrombosis disposition appears to be a markedly hypofibrinolytic response to venous occlusion of the paralysed lower extremities,^[139] a poor response likely attributable to low blood flow conditions in the paralysed lower extremities^[54,56] or interruption of adrenergic pathways that regulate fibrinolysis in the intact neuraxis.^[59]

2.6 Autonomic Dysregulation

Like those of the motor system, sympathetic nerve tracts descend in the spinal cord, albeit within the intermediolateral columns and not the corticospinal tracts, and exit with motor nerves in the thoracolumbar segments. As no sympathetic autonomic tracts exit the cord above the T₁ spinal level, individuals with cervical injuries often sustain decentralisation of their sympathetic nervous system. Loss of autonomic outflow to the adrenals is also observed in persons with paraplegia above the T₆ spinal level. Injury above the sacral cord segments abolishes central parasympathetic regulation of genitourinary organs (S2-4), which explains the common occurrences of neurogenic bowel and bladder after SCI, as well as erectile dysfunction in males.^[140] Autonomic dysfunction that results from injury above the thoracolumbar levels of sympathetic nerve outflow is associated with cardiac and circulatory dysfunction,^[34,132] clotting disorders,^[58] altered insulin metabolism,^[112,125] resting and exercise immunodysfunction,^[65,141-151] orthostatic incompetence,^[152] osteoporosis and joint deterioration,^[60,153] and thermal dysregulation at rest and during exercise.^[154-157] A blunted chronotropic response to exercise in persons with tetraplegia is well documented,^[158-160] and usually yields peak HRs in the mid-120 beats/min range, similar in magnitude to persons without SCI who exercise under conditions of pharmacological adrenergic blockade. The ab-

sence of, or meagre catecholamine responses to, exercise in these individuals have been reported^[161] and explains their suppressed HR responses to exercise. Thus, cardiac acceleration during exercise implies withdrawal of vagal tone and not increased sympathetic drive.

Few direct clinical comparisons can be made between persons having SCI above and below the highest level of sympathetic outflow at T₁ (i.e. between those persons whose motor status results in tetraplegia and paraplegia). Testing of subjects having inhomogeneous lesion levels usually leads to widely variable chronotropic, pressor, fuel, peripheral circulatory, thermal and work capacity responses. For those with paraplegia from T₂ to T₅ (or T₆), sparing of sympathetic efferents to the heart with resulting noradrenergic-mediated cardiac acceleration will be observed. A relatively normal response is observed below T₅ or T₆,^[162] as central inhibitory control of the adrenals (normally innervated from T₆-T₉) is regained below these levels.^[158]

3. Volitional Upper Extremity Exercise in Persons with SCI

3.1 Acute Physiological Responses to Exercise Stress

SCI blunt the expected physiological responses to exercise and diminish peak exercise capacity.^[163] These limitations are associated with the level of SCI, and are explained by various factors. First, ascending levels of injury cause greater loss of muscle mass necessary to serve as prime movers and stabilisers of trunk position. This requires that the arms simultaneously generate propulsive forces and steady the trunk during exercise. Secondly, increasing levels of injury are associated with increasing adrenergic dysfunction, and at key spinal levels dissociate adrenal, cardiac and total sympathetic nervous system control from the brain. As the adrenergic and noradrenergic systems play key roles in cardiovascular regulation and fuel homeostasis during exercise, their loss attenuates the cardiovascular

and metabolic efficiencies achieved in persons with an intact neuraxis.

Many studies have observed a direct association among level of injury, peak workload attained and peak oxygen uptake ($\dot{V}O_{2\text{peak}}$) reached during arm crank testing. This association serves as the basis for sports classification used during paralympic and other sports events for persons competing with physical disability. Exercise performance after SCI is limited by circulatory dysregulation accompanying thoracic injuries.^[138,156,159,164-166] Individuals with injuries below the level of sympathetic outflow at T₆ have a significantly lower resting SV and higher resting HR than persons without paraplegia.^[156,159,167] The significant elevation of resting and exercise HR is thought to compensate for a lower cardiac SV imposed by pooling of blood in the lower extremity venous circuits, diminished venous return and cardiac end-diastolic volumes, or frank circulatory insufficiency.^[138,168] Adjustments to the adrenergic systems after SCI may also regulate the excessive chronotropic response to work, as higher resting catecholamine levels and exaggerated catecholamine responses to physical work have been reported in individuals with paraplegia with middle thoracic (T₅) cord injuries. These exceed resting and exercise levels of both high level paraplegics and healthy persons without SCI.^[155,169] Hypersensitivity of the supraspinal spinal cord is believed to regulate this atypical adrenergic state and dynamic, which contrasts the downregulation of adrenergic functions observed in persons with high thoracic and cervical cord lesions.^[155,161]

An exaggerated HR response is experienced during physical activity by persons with paraplegia,^[158] which may be performance delimiting. This belief is consistent with reports in which subjects with paraplegia require higher levels of $\dot{V}O_2$ to perform the same work intensity as subjects without SCI^[32,138,156,170] and may represent a limiting factor in the performance of activities of daily living.^[21,156,171] As the sympathetic nervous system regulates haemodynamic and metabolic changes accompanying exercise, the elevated $\dot{V}O_2$ and HR response to work in individuals with paraplegia with

injuries below T₅ may be due to adrenergic overactivity accompanying their paraplegia.^[158,161,169]

3.2 Cardiorespiratory Testing

Assessment of cardiorespiratory fitness in persons with SCI requires specialised knowledge of both exercise testing procedures and the unique physiology that ensues SCI. Exercise testing modalities for these individuals commonly employ over-ground wheelchair propulsion on a treadmill or wheelchair rollers, cyclical arm ergometry and variations of electrically stimulated exercise. Exercise testing by wheelchair propulsion offers the advantage of comparison to many wheelchair-based daily tasks. However, wheelchair propulsion testing generally requires specialised equipment. Also, it is not known whether testing is best when using a single standardised wheelchair, in which the usual wheelchair-used interface is compromised, or having subjects use their habitual chair, in which some individuals may be advantaged by better equipment. These factors limit use of wheelchair ergometry as a research and clinical testing tool.

The exercise test mode most commonly used in the cardiorespiratory testing of persons with SCI is the arm crank ergometer. The lack of task specificity to wheelchair mobility has been considered a limitation to their usefulness. However, as these devices are generally found in the rehabilitation environment and allow standardised application of testing stresses, it represents the most frequently used means for clinical or fitness assessment of wheelchair users. Reliable test results require consistent positioning of the test subject and the ergometer. The test subject and the cranking ergometer should be positioned so that the axis of the crank arm is horizontally aligned with the subject's shoulder joint and the elbows are slightly flexed at the point of furthest reach. To ensure enhanced reliability of test results, care should be taken to replicate subject positioning and to use the same wheelchair and cushion. The latter may affect seating stability and exercise efficiency.

Cardiorespiratory exercise testing of persons with SCI follows the same general protocols as

testing in the general population. In most cases, the graded multi-stage tests proceed in 2- to 3-minute stages until volitional exhaustion. As $\dot{V}O_{2peak}$ is dependent on level of SCI, lower intensity loads are applied than those used in standard exercise testing. These workloads have been detailed in several publications,^[172-175] although no particular advantage for a single testing strategy has been tested or established. However, accurate and valid results require that testing utilise metabolic monitoring to establish the $\dot{V}O_{2peak}$, as prediction equations normally used to estimate maximal responses in persons without disability have not been validated in those with SCI.

3.3 Endurance Training

In most cases, injury to the spinal cord renders the lower extremities without sufficient strength, endurance or motor control to support safe and effective physical training. This explains why most exercise training after SCI employs upper extremity exercise modes including arm crank ergometry, wheelchair ergometry and swimming. All of these training modes improve $\dot{V}O_{2peak}$ in those with SCI,^[27,32,172,176-186] with the magnitude of improvement inversely proportional to level of spinal lesion. While it is possible for persons with low tetraplegia to train on an arm ergometer, special measures must be taken to affix the hands to an ergometer and their gains in $\dot{V}O_{2peak}$ fail to match those of their SCI counterparts with paraplegia.^[187] Therefore, the level of injury is a key to predicting outcome from arm endurance training.^[188,189] Guidelines for training after SCI have been published by several authorities,^[32,52,172] although testing for superiority of training algorithms has not been performed. However, given that most persons with SCI are sedentary long after injury and the accepted dictum that training best benefits the most deconditioned individuals, these methods are nonetheless successful in achieving higher levels of fitness.

Benefits of arm conditioning are widely reported for persons with SCI, with many studies reporting significant increases in $\dot{V}O_{2peak}$ after training.^[27,185-187,190,191] It is less clear whether the changes in $\dot{V}O_{2peak}$ result from central (HR, SV,

cardiac output [\dot{Q}]) adaptations or increases in peripheral oxygen extraction ($a-\dot{V}O_2$ difference), or whether central adaptations take place at all. Only one endurance training study examining 16 weeks of intense arm exercise observed clear evidence of central cardiovascular benefit.^[176] Notwithstanding the mechanisms contributing to increased $\dot{V}O_{2peak}$ after arm exercise training, many studies report individuals with paraplegia improve their $\dot{V}O_{2peak}$ by 10–20% following 8–12 weeks of training. These gains are inversely proportional to the level of SCI, as those with higher levels of injury attain lower peak exercise capacities on arm exercise test than those with lower-level SCI, and both groups lower than persons without disability tested by arm ergometry.

Endurance training recommendations for those with SCI do not vary dramatically from advice directed to the general population.^[192-194] The American College of Sports Medicine recommended training frequencies, durations and intensities are contained in *Exercise Management for Persons with Chronic Diseases and Disabilities*.^[192] Generally, three to five weekly exercise sessions of 20–60 minutes in duration and at an intensity of 50–80% $\dot{V}O_{2peak}$ is the recommended exercise prescription for persons with paraplegia using the following modes of exercise:

- arm cranking
- wheelchair propulsion
- swimming
- wheelchair sports
- circuit resistance training (CRT)
- electrically stimulated cycling
- electrically stimulated walking.

This reference suggests that target exercise intensities should produce HR responses equivalent to 50–80% of the individual's peak HR. As in the general population, excessive exercise volume has been associated with increase pain and injury. Thus, exercise frequency and duration should be carefully monitored and gradually increased with the input of certified exercise professionals familiar with the unique physiological responses of persons with SCI to exercise. As long-term compliance and injury

avoidance are major goals of training, erring on the conservative side of selected exercise durations and intensities are prudent and more important for persons training with a disability than those without.

3.4 Resistance Training for Persons with SCI

Far less is known about effects of resistance than endurance training in persons with SCI. This might seem counterintuitive, as muscle weakness has been reported to precipitate pain and dysfunction as persons with SCI age with their disability. Moreover, the most commonly reported symptom of upper extremity physical dysfunction among persons with SCI is pain of the shoulder joint and girdle.^[66,195-200] While a single cause for shoulder pain has not been identified, deterioration and injury resulting from insufficient shoulder strength is one commonly cited source.^[63,199-202] Pain that accompanies wheelchair locomotion and other wheelchair activities reportedly interferes with functional activities including upper extremity weight bearing for transfers, high resistance muscular activity in extremes of range of motion, wheelchair propulsion up inclines and frequent overhead activity.^[196,198-200] Onset of pain is common during body transfer activities and severity increases as time following injury lengthens.^[196] Given evidence that wheelchair locomotion is a major source of pain and dysfunction for persons with SCI, incorporation of resistance training into the healthcare plan of those with SCI appears both justified and essential.

Several investigators have studied effects of resistance exercise in persons with paraplegia. In a study of Scandinavian men (most of whom had incomplete low thoracic lesions) a weight-training programme emphasising triceps strengthening for crutch walking was undertaken for 7 weeks with modest but significant increases in $\dot{V}O_{2\text{peak}}$ observed following training. These endurance gains were accompanied by increased strength of the triceps brachii.^[203] Another study^[204] examined the effects of arm ergometry in subjects assigned to high-intensity (70% of their $\dot{V}O_{2\text{peak}}$) or low-intensity training (40% of their $\dot{V}O_{2\text{peak}}$) for 20 or 40 minutes per session, respectively. Strength gains

were limited to subjects assigned high-intensity training, and occurred only in the shoulder joint extensor and elbow flexor muscles. Otherwise, no changes in shoulder joint abduction or adduction strengths were reported, and none of the muscles that move or stabilise the scapulothoracic articulation or chest were stronger following training. These results suggest that arm crank exercise is ineffective as a training mode for upper extremity strengthening because it fails to target the muscles most involved in performance of daily activities.

Conditioning of five paraplegic and five tetraplegic subjects three times weekly for 9 weeks was reported using a hydraulic fitness machine.^[205] Exercises were limited to two manoeuvres: (i) chest press/chest row; and (ii) shoulder press/latisimus pull.^[205] Significant increases in peak values of $\dot{V}O_2$ and power output measured by arm ergometry testing were observed at the conclusion of the study, although no testing was conducted that directly measured strength gain in any muscle groups undergoing training. Another study focused their training on strengthening of the scapular muscles, although this study focused solely on the scapular retractor muscles when comparing seated rowing and a standardised scapular retraction exercise, and did so only for concentric actions.^[206] The authors found that higher levels of retractor activation were obtained during backward compared with forward wheelchair locomotion and suggested that rowing was effective for improving scapular retractor activity and cardiorespiratory fitness. A recent study observed reduced shoulder pain following a series of shoulder resistance exercises using elastic bands.^[63]

As both endurance and resistance exercise benefit those without SCI, the effects of CRT^[207] on various attributes of fitness, dyslipidaemia and shoulder pain have been studied in young and middle-aged subjects with paraplegia. This exercise programme incorporates periods of low-intensity high-paced movements interposed within activities performed at a series of resistance training stations. The CRT exercise programme adapted for persons with paraplegia consisted of three circuits of six resistance stations encompassing three pairs of agonist/

antagonist movements (e.g. overhead press and pull) and three 2-minute periods of free wheeling arm cranking performed between resistance manoeuvres. No true rest periods were allowed during the performance of CRT, with active recovery limited to the time necessary for the subject to propel the wheelchair to the next exercise station. Three weekly sessions were completed with each session lasting approximately 45 minutes. Young subjects undergoing 16 weeks of mixed resistance and endurance exercise increased their arm $\dot{V}O_{2peak}$ by an average of 29%, with accompanying strength gains of 13–40%, depending on the upper extremity site tested.^[174] Subjects undergoing CRT lowered their total and LDL-C while increasing their HDL-C by nearly 10%.^[175] Subjects aged over 40 years undergoing the same treatment for 12 weeks experienced significant gains in all of endurance, strength and anaerobic power, even though the latter was not specifically targeted by training (Nash MS, unpublished data). Shoulder pain present in these subjects before training was assessed by a pain instrument validated in the population and significantly reduced. Pain was eliminated in four of ten subjects.

In a separate study, the effects of circuit training performed on an adapted multi-exercise system could be replicated by use of elastic bands.^[208] Evidence therefore supports health and fitness advantages of CRT over either endurance or resistance exercises alone for persons with paraplegia.

General recommendations for strength training exercise prescription include three sets of 8–12 repetitions per exercise movement for two sessions per week at moderate to high intensity using the following modes of exercise:

- free weights
- weight machines
- elastic tubing and bands.

4. Electrically Stimulated Exercise in Persons with SCI

4.1 Basis of Electrically Stimulated Movement

Many forms of electrically stimulated exercise are available for use by persons with SCI. These

include site-specific stimulation of the lower extremities^[82,209-212] and upper extremities,^[65,213-218] leg cycling,^[30,34,134,219-222] leg exercise with upper extremity assist,^[222-225] lower body rowing,^[50] electrically assisted arm ergometry,^[226,227] electrically stimulated standing^[76,217] and electrically stimulated bipedal ambulation either with^[74-76,228,229] or without an orthosis.^[55,70,72,173,230-232] Most forms of exercise require that the lower motor neuron system remain intact following injury, as direct activation of the peripheral nerve and not the muscle by electrical current ultimately results in muscle activation.^[233] This reality excludes most individuals having cauda equina or conus medullaris syndromes from electrically stimulated exercise. It may also compromise the efficiency of a muscle having a denervated segment from injury to its anterior horn cells or spinal (Wallnerian) degeneration from injured adjacent spinal areas. Many applications of SCI target muscle strengthening of limb segments whose motor function is partially spared by injury,^[76] while others use electrical current as a neuroprosthesis for the lower extremities^[71,229] and upper extremities.^[234-238]

Most electrical stimulation devices currently approved for use by the US FDA employ surface, not implanted, electrical stimulation. The most common uses of surface electrical stimulation for system exercise in those with SCI include electrically stimulated cycling, either with or without upper extremity assistive propulsion and electrically stimulated ambulation. Qualifications necessary to safely participate in exercise programmes have been described^[71,72,233,239,240] and risks of participation are included in the section of this monograph addressing adverse events associated with exercise by those with SCI.

4.2 Leg Cycling Exercise

The simplest form of multi-limb segmental exercise for persons with SCI employs a cycling motion performed in a slightly recumbent seated position. This electrically stimulated cycling uses electrically activated contractions of the bilateral quadriceps, hamstrings and gluteus muscles under computer microprocessor command.^[31] Control of pedal ca-

dence and muscle stimulation intensity is exerted by feedback provided from position sensors placed in the pedal gear.^[241]

Training with electrically stimulated cycling is often preceded with electrically stimulated strengthening of the quadriceps muscles, which is necessary in cases of severe muscle atrophy or diminished muscle endurance.^[209,221] Responses to electrically stimulated exercise are variable, ranging from poor to robust. As noted, poor muscle strength and endurance and altered muscle contractile properties normally accompany SCI. In general, these factors will slow success in training, especially in those individuals with longstanding paralysis, absent spasticity and flexor patters of spasticity. Electrically stimulated strength training programmes are capable of altering the strength, resistance to fatigue and contractile properties of the muscles undergoing the training. Dudley and associates,^[209] demonstrated the effectiveness of a simple programme of electrically induced knee extensions, performed twice weekly over an 8-week period, dramatically reversed muscular atrophy of the quadriceps muscles.

Notwithstanding poor levels of muscle strength and endurance early in most training programmes and despite limited ability to exercise against intense workloads, enhanced levels of fitness,^[224,242,243] improved gas exchange kinetics^[244,245] and increased muscle mass^[218] have been reported following exercise training using electrically stimulated cycling. For those with neurologically incomplete injuries, gains in lower extremity mass, as well as isometric strength and endurance under conditions of voluntary and electrically stimulated cycling have been reported.^[218] Reversal of the adaptive left ventricular atrophy reported in persons with tetraplegia has also been observed, with near normalisation of pre-training cardiac mass.^[134] This change may be associated with significantly improved lower extremity circulation following training,^[246,247] which is also accompanied by a more robust hyperaemic response to experimental occlusion ischaemia.^[55,56]

Attenuation of paralytic osteopaenia has been observed by several investigators^[248,249] and an in-

creased rate of bone turnover by another,^[10] although the sites benefiting from training are the lumbar spine and proximal tibia, not the proximal femur.^[249] Not all studies have found a post-training increase in mineral density for bones located below the level of the lesion.^[250] Those that fail to do so have usually studied subjects with longstanding paralysis, in which attenuation or reversal of osteopaenia by any treatment has yet to be reported. Notwithstanding, a study examining the appearance of lower extremity joints and joint surfaces using magnetic resonance imaging reported no degenerative changes induced by cycling, and less joint surface necrosis than previously reported in sedentary persons with SCI.^[60] Improved body composition favouring increased lean mass and decreased fat mass^[251] and an enhancement of whole-body insulin uptake, insulin-stimulated 3-O-methyl glucose transport and increased expression of GLUT4 transport protein in the quadriceps muscle have been reported.^[252] In a recent report, this training has resulted in improved profiles of insulin resistance.^[253] When combined with simultaneous upper extremity arm ergometry, the acute cardiovascular metabolic responses to electrically stimulated cycling are more intense and the gains in fitness greater than observed with lower extremity cycling alone.^[224,253]

4.3 Bipedal Ambulation

Complex electrical stimulation to achieve bipedal ambulation has been used as a neuroprosthesis for those with motor-complete injuries^[71,72,228,239,254] and an assistive neuroprosthesis for persons with incomplete SCI who lack necessary strength to support independent ambulation.^[255-263] Surface and implantable neuroprostheses for those without spared motor function have both been fabricated,^[76,264] although the sole method currently approved by the FDA (Parastep-1®)¹ uses surface electrical stimulation of the quadriceps and gluteus muscles.^[72,230] Muscle activation is sequenced by a microprocessor worn on the belt, with activation of step initiated by a finger-sensitive control switch located on a rolling

1 The use of trade names is for product identification purposes only and does not imply endorsement.

walker used by ambulating subjects. When pressed, the electrical stimulator sends a current to the stance limb that initiates contraction of the quadriceps and gluteus muscles. Contralateral hip flexion is then achieved by exploiting an ipsilateral flexor withdrawal reflex obtained by introducing a nociceptive electrical stimulus over the common peroneal nerve at the head of the fibula. This allows the hip, knee and ankle to move into flexion followed by extension of the knee joint by electrical stimulation to the quadriceps. As muscle fatigue occurs increasing levels of stimulation can be provided by a switch mounted on the handle of the rolling walker.

As in the case of electrically stimulated cycling, functional use of ambulation neuroprostheses by those with paraplegia are compromised by post-injury muscle weakness and poor endurance. These need to be addressed in subject preparation before ambulation training is initiated. Electrically activated knee extensions against progressively increased resistance (sandbag at ankle) is the standard training exercise. While not as commonly addressed, conditioning of the gluteal muscles with electrically produced hip bridging (in supine position) prior to bodyweight-supported gait training will enhance the response of the hip extensors to electrical stimulation. A single most effective protocol to satisfy requisite strength and endurance needs for upright ambulation training has not been reported. However, it is inadvisable to undertake electrically stimulated ambulation training until the ability to stand with electrical stimulation for at least 3–5 minutes has been demonstrated.

Once training is initiated, rates of ambulation are relatively slow and distances of ambulation relatively limited when using the system,^[229] and the community use of these devices remains limited to a small percentage of training subjects. Despite the limitations of ambulation velocity and distance, ambulation distances of up to 1.6km (1 mile) have been reported after training in some subjects^[229] and upper extremity fitness is enhanced following ambulation training.^[71,173] Other adaptations to training include significantly increased lower extremity muscle mass,^[265] improved resting blood flow^[55] and an

augmented hyperaemic response to an experimental ischaemic stimulus.^[55] No change in lower extremity bone mineralisation following has been reported,^[232] although most subjects participating in research trials of electrically stimulated exercise are beyond the duration of injury at which reversal of their neurogenic osteoporosis is expected.^[232]

5. Risks of Exercise When Performed by Persons with SCI

Special attention is required when designing, instituting or performing exercise programmes for persons with SCI. Some of the risks encountered will be similar to those experienced by persons without paralysis, although complications such as general overuse may be exaggerated in persons with SCI, and their occurrence will likely compromise daily activities to a far greater extent than similar injuries arising in persons without SCI.

5.1 Autonomic Dysreflexia

Individuals having cord injuries at or above the T6 spinal level are prone to episodes of autonomic hyperreflexia when exposed to noxious stimuli.^[266] The neurological basis for these episodes involves loss of supraspinal sympathetic inhibition that normally suppresses the unrestricted autonomic reflex accompanying such exposure. This allows the adrenals to release high concentrations of epinephrine (adrenaline) under reflex control and infralesional adrenergic targets to experience the full measure of reflex noradrenergic stimulation.^[267] The most common stimuli evoking autonomic dysreflexia are bladder and bowel distention before their emptying. Other stimuli include venous thromboembolism, bone fracture, sudden temperature change, febrile episodes and exercise. The disposition to autonomic dysreflexia during exercise is especially heightened when an electrical current is used to generate muscle movement, or when exercising while febrile or during bladder emptying. Episodes of autonomic dysreflexia are characterised by hypertension and bradycardia, supralesional erythema, piloerection and headache. In some cases, hypertension can rise to the point where crisis headache results, and cere-

bral haemorrhage and death might ensue. Recognition of these episodes, withdrawal of the offending stimulus, and the possible administration of a fast-acting peripheral vasodilator may be critical in preventing serious medical complications. Prophylaxis with a slow calcium channel antagonist or α_1 -selective adrenergic antagonist may be needed prior to exercise.^[268-270] It is known that wheelchair racers have intentionally induced dysreflexia as an ergogenic aid by restricting urine outflow through a Foley catheter,^[271] which represents a dangerous and possibly life-threatening practice.

5.2 Musculoskeletal Injury

Fracture and joint dislocation of the lower extremities is a risk of participation in exercise by those with SCI, and may be caused by asynergistic movement of limbs against the force imposed by either electrical stimulation or the device used for exercise.^[36] This explains why these activities are contraindicated for individuals with severe spasticity or spastic response to the introduction of electrical current. Precautions to prevent overuse injuries of the arms and shoulders must be taken for those participating in upper extremity exercise.^[201,206,272] As the shoulder joints are mechanically ill-suited to perform locomotor activities, but must do so in individuals using a manual wheelchair for transportation, these injuries may ultimately compromise performance of essential daily activities including wheelchair propulsion, weight relief and depression transfers.^[273,274]

5.3 Hypotension

Small risks of post-exercise hypotension^[131,275] are associated with lost vasomotor responses to orthostatic repositioning,^[276] although these episodes can abate after upper limb training.

5.4 Thermal Dysregulation

Individuals with SCI often lack sudomotor responses below their level of injury and are thus challenged to maintain thermal stability.^[154,277-279] These responses are less pronounced as the level of SCI descends,^[280,281] and when exercising in an en-

vironment controlled for temperature and humidity.^[278,282] Thus, attention should be paid to hydration and, if possible, limiting the duration and intensity of activities performed in intemperate environments.

6. Conclusions

Multisystem organ dysfunction is common after SCI. Depending on the level of injury, alterations of cardiac function, peripheral circulation, autonomic function, skeletal integrity, muscle composition and genitourinary functions all accompany an SCI. The fact that many of these pathological system changes are reversible permits better understanding of the roles played by central innervation and physical activity on organ system function in persons with intact neuraxes. As levels of injury, types of injury and the extent of organ system dysfunction vary among persons with SCI, careful attention must be paid to accurate classification of individuals before entry into clinical treatment or study. The use of homogeneous subject populations based on similar levels, types and durations of injury is especially important if conclusions concerning deconditioning and reconditioning in these persons are to be reached.

Despite these special needs and warnings, many persons with SCI benefit from sustained therapeutic and recreational exercise. Individuals with higher levels of cord injury may require electrical stimulation to perform exercise, which poses special restrictions on use and unique risks from participation. Qualified individuals safely improved cardiovascular and musculoskeletal functions. Positive benefits of training on bone density, regulation of orthostatic tolerance and affect have been reported in studies with limited numbers of subjects and require well controlled investigations for confirmation. Individuals with spared motor control of the upper extremity can perform arm or wheelchair exercises and participate in recreational sports. Greater emphasis needs to be placed on strengthening of the upper extremity to preserve shoulder and arm functions for performance of daily living as these individuals age with their paralysis. Risks of injury or illnesses associat-

ed with imprudent exercise must be managed to ensure that the desirable benefits of physical activity can be sustained. If carefully prescribed, exercise has the demonstrated ability to enhance the activity, life satisfaction and health of those with disability from SCI.

Acknowledgements

The preparation of this manuscript was made possible by funding from the Miami Project to Cure Paralysis, University of Miami, Florida, USA, and the Rehabilitation Research and Development Service, Department of Veterans Affairs, USA. The authors have no conflicts of interest directly relevant to the content of this review.

References

- DeVivo MJ, Black KJ, Stover SL. Causes of death during the first 12 years after spinal cord injury. *Arch Phys Med Rehabil* 1993; 74: 248-54
- DeVivo MJ, Go BK, Jackson AB. Overview of the national spinal cord injury statistical center database. *J Spinal Cord Med* 2002; 25: 335-8
- DeVivo MJ, Shewchuk RM, Stover SL, et al. A cross-sectional study of the relationship between age and current health status for persons with spinal cord injuries. *Paraplegia* 1992; 30: 820-7
- Ergas Z. Spinal cord injury in the United States: a statistical update. *Cent Nerv Syst Trauma* 1985; 2: 19-32
- Nash MS. Cardiovascular fitness after spinal cord injuries. In: Lin V, editor. *Spinal cord medicine: principles and practice*. New York (NY): Demos Medical Publications, 2002: 637-46
- Nash MS, Horton JA. Recreational and therapeutic exercise after SCI. In: Kirshbaum S, Campagnolo DI, DeLisa JS, editors. *Spinal cord medicine*. Philadelphia (PA): Lippincott, Williams and Wilkins, 2002: 331-7
- Dearwater SR, LaPorte RE, Robertson RJ, et al. Activity in the spinal cord-injured patient: an epidemiologic analysis of metabolic parameters. *Med Sci Sports Exerc* 1986; 18: 541-4
- Noreau L, Shephard RJ, Simard C, et al. Relationship of impairment and functional ability to habitual activity and fitness following spinal cord injury. *Int J Rehabil Res* 1993; 16: 265-75
- Washburn RA, Figoni SF. Physical activity and chronic cardiovascular disease prevention in spinal cord injury: a comprehensive literature review. *Top Spinal Cord Injury Rehabil* 1998; 3: 16-32
- Bloomfield SA, Mysiw WJ, Jackson RD. Bone mass and endocrine adaptations to training in spinal cord injured individuals. *Bone* 1996; 19: 61-8
- Bostom AG, Toner MM, McArdle WD, et al. Lipid and lipoprotein profiles relate to peak aerobic power in spinal cord injured men. *Med Sci Sports Exerc* 1991; 23: 409-14
- Demirel G, Yilmaz H, Paker N, et al. Osteoporosis after spinal cord injury. *Spinal Cord* 1998; 36: 822-5
- Hjeltnes N, Jansen T. Physical endurance capacity, functional status and medical complications in spinal cord injured subjects with long-standing lesions. *Paraplegia* 1990; 28: 428-32
- Kocina P. Body composition of spinal cord injured adults. *Sports Med* 1997; 23: 48-60
- Naftchi NE, Viau AT, Sell GH, et al. Mineral metabolism in spinal cord injury. *Arch Phys Med Rehabil* 1980; 61: 139-42
- Nash MS. Immune responses to nervous system decentralization and exercise in quadriplegia. *Med Sci Sports Exerc* 1994; 26: 164-71
- Nash MS. Exercise reconditioning of the heart and peripheral circulation after spinal cord injury. *Top Spinal Cord Inj Rehabil* 1997; 3: 1-15
- Ragnarsson KT. The cardiovascular system. In: Whiteneck GG, Carlifue SW, Gerhart KA, et al., editors. *Aging with spinal cord injury*. New York: Demos Publications, 1993: 73-92
- Shields RK. Muscular, skeletal, and neural adaptations following spinal cord injury. *J Orthop Sports Phys Ther* 2002; 32: 65-74
- Dallmeijer AJ. Spinal cord injury and physical activity: wheelchair performance in rehabilitation and sports [dissertation]. Amsterdam: Vrije Universiteit (Free University), Faculty of Human Movement Sciences, 1998
- Janssen TW, van Oers CA, Veeger HE, et al. Relationship between physical strain during standardised ADL tasks and physical capacity in men with spinal cord injuries. *Paraplegia* 1994; 32: 844-59
- Bauman WA, Spungen AM. Disorders of carbohydrate and lipid metabolism in veterans with paraplegia or quadriplegia: a model of premature aging. *Metabolism* 1994; 43: 749-56
- Bauman WA, Spungen AM, Adkins RH, et al. Metabolic and endocrine changes in persons aging with spinal cord injury. *Assist Technol* 1999; 11: 88-96
- Gerhart KA, Bergstrom E, Charlifue SW, et al. Long-term spinal cord injury: functional changes over time. *Arch Phys Med Rehabil* 1993; 74: 1030-4
- Ohry A, Shemesh Y, Rozin R. Are chronic spinal cord injured patients (SCIP) prone to premature aging? *Med Hypotheses* 1983; 11: 467-9
- Whiteneck G, Menter RR. Where do we go from here? In: Whiteneck GG, Cahlifue SW, Gerhart KA, et al., editors. *Aging with spinal cord injury*. New York: Demos, 1993: 361-9
- Cowell LL, Squires WG, Raven PB. Benefits of aerobic exercise for the paraplegic: a brief review. *Med Sci Sports Exerc* 1986; 18: 501-8
- Davis GM. Exercise capacity of individuals with paraplegia. *Med Sci Sports Exerc* 1993; 25: 423-32
- Figoni SF. Perspectives on cardiovascular fitness and SCI. *J Am Paraplegia Soc* 1990; 13: 63-71
- Glaser RM. Exercise and locomotion for the spinal cord injured. *Exerc Sport Sci Rev* 1985; 13: 263-303
- Glaser RM. Functional neuromuscular stimulation: exercise conditioning of spinal cord injured patients. *Int J Sports Med* 1994; 15: 142-8
- Hoffman MD. Cardiorespiratory fitness and training in quadriplegics and paraplegics. *Sports Med* 1986; 3: 312-30
- Jacobs PL, Nash MS. Modes, benefits, and risks of voluntary and electrically induced exercise in persons with spinal cord injury. *J Spinal Cord Med* 2001; 24: 10-8
- Nash MS, Bilsker MS, Kearney HM, et al. Effects of electrically-stimulated exercise and passive motion on echocardiographically-derived wall motion and cardiodynamic function in tetraplegic persons. *Paraplegia* 1995; 33: 80-9
- Thomas AJ, Davis GM, Sutton JR. Cardiovascular and metabolic responses to electrical stimulation-induced leg exercise in spinal cord injury. *Methods Inf Med* 1997; 36: 372-5

36. Hartkopp A, Murphy RJ, Mohr T, et al. Bone fracture during electrical stimulation of the quadriceps in a spinal cord injured subject. *Arch Phys Med Rehabil* 1998; 79: 1133-6
37. Bracken MB, Shepard MJ, Collins WF, et al. A randomized, controlled trial of methylprednisolone or naloxone in the treatment of acute spinal-cord injury. Results of the Second National Acute Spinal Cord Injury Study. *N Engl J Med* 1990; 322: 1405-11
38. Ditunno Jr JF. The John Stanley Coulter Lecture. Predicting recovery after spinal cord injury: a rehabilitation imperative. *Arch Phys Med Rehabil* 1999; 80: 361-4
39. Marino RJ, Ditunno Jr JF, Donovan WH, et al. Neurologic recovery after traumatic spinal cord injury: data from the Model Spinal Cord Injury Systems. *Arch Phys Med Rehabil* 1999; 80: 1391-6
40. Ditunno Jr JF, Graziani V, Tessler A. Neurological assessment in spinal cord injury. *Adv Neurol* 1997; 72: 325-33
41. American Spinal Injury Association. International standards for neurological classification of spinal cord injury, revised 2002. Chicago (IL): American Spinal Injury Association, 2002
42. Marino RJ, Ditunno Jr JF, Donovan WH, et al. Neurologic recovery after traumatic spinal cord injury: data from the Model Spinal Cord Injury Systems. *Arch Phys Med Rehabil* 1999; 80: 1391-6
43. Maynard Jr FM, Bracken MB, Creasey G, et al. International standards for neurological and functional classification of spinal cord injury. American Spinal Injury Association. *Spinal Cord* 1997; 35: 266-74
44. Frankel HL, Hancock DO, Hyslop G, et al. The value of postural reduction in the initial management of closed injuries of the spine with paraplegia and tetraplegia. I. Paraplegia 1969; 7: 179-92
45. Waters RL, Adkins RH, Yakura JS. Definition of complete spinal cord injury. *Paraplegia* 1991; 29: 573-81
46. Ko HY, Ditunno Jr JF, Graziani V, et al. The pattern of reflex recovery during spinal shock. *Spinal Cord* 1999; 37: 402-9
47. Little JW, Ditunno Jr JF, Stiens SA, et al. Incomplete spinal cord injury: neuronal mechanisms of motor recovery and hyperreflexia. *Arch Phys Med Rehabil* 1999; 80: 587-99
48. Nacimiento W, Noth J. What, if anything, is spinal shock? *Arch Neurol* 1999; 56: 1033-5
49. White RJ, Likavec MJ. Spinal shock: spinal man. *J Trauma* 1999; 46: 979-80
50. Laskin JJ, Ashley EA, Olenik LM, et al. Electrical stimulation-assisted rowing exercise in spinal cord injured people: a pilot study. *Paraplegia* 1993; 31: 534-41
51. Bauman WA, Spungen AM. Metabolic changes in persons after spinal cord injury. *Phys Med Rehabil Clin N Am* 2000; 11: 109-40
52. Phillips WT, Kiratli BJ, Sarkarati M, et al. Effect of spinal cord injury on the heart and cardiovascular fitness. *Curr Probl Cardiol* 1998; 23: 641-716
53. Hopman MT, Monroe M, Dueck C, et al. Blood redistribution and circulatory responses to submaximal arm exercise in persons with spinal cord injury. *Scand J Rehabil Med* 1998; 30: 167-74
54. Jacobs PL, Mahoney ET, Robbins A, et al. Hypokinetic circulation in persons with paraplegia. *Med Sci Sports Exerc* 2002; 34: 1401-7
55. Nash MS, Jacobs PL, Montalvo BM, et al. Evaluation of a training program for persons with SCI paraplegia using the Parastep 1 ambulation system: part 5: lower extremity blood flow and hyperemic responses to occlusion are augmented by ambulation training. *Arch Phys Med Rehabil* 1997; 78: 808-14
56. Nash MS, Montalvo BM, Applegate B. Lower extremity blood flow and responses to occlusion ischemia differ in exercise-trained and sedentary tetraplegic persons. *Arch Phys Med Rehabil* 1996; 77: 1260-5
57. Webb P, Saris WH, Schoffelen PF, et al. The work of walking: a calorimetric study. *Med Sci Sports Exerc* 1988; 20: 331-7
58. Green D, Hull RD, Mammen EF, et al. Deep vein thrombosis in spinal cord injury: summary and recommendations. *Chest* 1992; 102: 633S-5S
59. Winther K, Gleerup G, Snorrason K, et al. Platelet function and fibrinolytic activity in cervical spinal cord injured patients. *Thromb Res* 1992; 65: 469-74
60. Nash MS, Tehranzadeh J, Green BA, et al. Magnetic resonance imaging of osteonecrosis and osteoarthritis in exercising quadriplegics and paraplegics. *Am J Phys Med Rehabil* 1994; 73: 184-92
61. Segatore M. The skeleton after spinal cord injury, part 1: theoretical aspects. *SCI Nurs* 1995; 12: 82-6
62. Segatore M. The skeleton after spinal cord injury, part 2: management of sublesional osteoporosis. *SCI Nurs* 1995; 12: 115-20
63. Curtis KA, Tyner TM, Zachary L, et al. Effect of a standard exercise protocol on shoulder pain in long-term wheelchair users. *Spinal Cord* 1999; 37: 421-9
64. Levi R, Hultling C, Nash MS, et al. The Stockholm spinal cord injury study. I. medical problems in a regional SCI population. *Paraplegia* 1995; 33: 308-15
65. Nash MS. The immune system. In: Whiteneck G, Charlifue SW, Gerhart KA, et al., editors. *Aging with spinal cord injury*. New York: Demos Publications, 1993: 159-82
66. Sie IH, Waters RL, Adkins RH, et al. Upper extremity pain in the postrehabilitation spinal cord injured patient. *Arch Phys Med Rehabil* 1992; 73: 44-8
67. Waring WP, Maynard FM. Shoulder pain in acute traumatic quadriplegia. *Paraplegia* 1991; 29: 37-42
68. LaPorte RE, Adams LL, Savage DD, et al. The spectrum of physical activity, cardiovascular disease and health: an epidemiologic perspective. *Am J Epidemiol* 1984; 120: 507-17
69. Curtis KA, McClanahan S, Hall KM, et al. Health, vocational, and functional status in spinal cord injured athletes and nonathletes. *Arch Phys Med Rehabil* 1986; 67: 862-5
70. Davis R, Houdayer T, Andrews B, et al. Paraplegia: prolonged closed-loop standing with implanted nucleus FES- 22 stimulator and Andrews' foot-ankle orthosis. *Stereotact Funct Neurosurg* 1997; 69: 281-7
71. Graupe D, Kohn KH. Functional neuromuscular stimulator for short-distance ambulation by certain thoracic-level spinal-cord-injured paraplegics. *Surg Neurol* 1998; 50: 202-7
72. Klose KJ, Jacobs PL, Broton JG, et al. Evaluation of a training program for persons with SCI paraplegia using the Parastep 1 ambulation system, part I: ambulation performance and anthropometric measures. *Arch Phys Med Rehabil* 1997; 78: 789-93
73. Marsolais EB, Edwards BG. Energy costs of walking and standing with functional neuromuscular stimulation and long leg braces. *Arch Phys Med Rehabil* 1988; 69: 243-9
74. Solomonow M, Reisin E, Aguilar E, et al. Reciprocating gait orthosis powered with electrical muscle stimulation (RGO II), part II: medical evaluation of 70 paraplegic patients. *Orthopedics* 1997; 20: 411-8

75. Thoumie P, Le Claire G, Beillot J, et al. Restoration of functional gait in paraplegic patients with the RGO-II hybrid orthosis: a multicenter controlled study, II. physiological evaluation. *Paraplegia* 1995; 33: 654-9
76. Triolo RJ, Bieri C, Uhlir J, et al. Implanted functional neuromuscular stimulation systems for individuals with cervical spinal cord injuries: clinical case reports. *Arch Phys Med Rehabil* 1996; 77: 1119-28
77. Bernard PL, Mercier J, Varray A, et al. Influence of lesion level on the cardioventilatory adaptations in paraplegic wheelchair athletes during muscular exercise. *Spinal Cord* 2000; 38: 16-25
78. Bauman WA, Spungen AM, Zhong YG, et al. Depressed serum high density lipoprotein cholesterol levels in veterans with spinal cord injury. *Paraplegia* 1992; 30: 697-703
79. Burnham R, Martin T, Stein R, et al. Skeletal muscle fibre type transformation following spinal cord injury. *Spinal Cord* 1997; 35: 86-91
80. Castro MJ, Apple Jr DF, Hillegass EA, et al. Influence of complete spinal cord injury on skeletal muscle cross-sectional area within the first 6 months of injury. *Eur J Appl Physiol Occup Physiol* 1999; 80: 373-8
81. Castro MJ, Apple Jr DF, Rogers S, et al. Influence of complete spinal cord injury on skeletal muscle mechanics within the first 6 months of injury. *Eur J Appl Physiol* 2000; 81: 128-31
82. Greve JM, Muszkat R, Schmidt B, et al. Functional electrical stimulation (FES): muscle histochemical analysis. *Paraplegia* 1993; 31: 764-70
83. Grimby G, Broberg C, Krotkiewska I, et al. Muscle fiber composition in patients with traumatic cord lesion. *Scand J Rehabil Med* 1976; 8: 37-42
84. Castro MJ, Apple Jr DF, Staron RS, et al. Influence of complete spinal cord injury on skeletal muscle within 6 mo of injury. *J Appl Physiol* 1999; 86: 350-8
85. Levy M, Mizrahi J, Susak Z. Recruitment, force and fatigue characteristics of quadriceps muscles of paraplegics isometrically activated by surface functional electrical stimulation. *J Biomed Eng* 1990; 12: 150-6
86. Rochester L, Chandler CS, Johnson MA, et al. Influence of electrical stimulation of the tibialis anterior muscle in paraplegic subjects, 1: contractile properties. *Paraplegia* 1995; 33: 437-49
87. Andersen JL, Mohr T, Biering-Sorensen F, et al. Myosin heavy chain isoform transformation in single fibres from m. vastus lateralis in spinal cord injured individuals: effects of long-term functional electrical stimulation (FES). *Pflugers Arch* 1996; 431: 513-8
88. Lotta S, Scelsi R, Alfonsi E, et al. Morphometric and neurophysiological analysis of skeletal muscle in paraplegic patients with traumatic cord lesion. *Paraplegia* 1991; 29: 247-52
89. Scelsi R, Marchetti C, Poggi P, et al. Muscle fiber type morphology and distribution in paraplegic patients with traumatic cord lesion: histochemical and ultrastructural aspects of rectus femoris muscle. *Acta Neuropathol (Berl)* 1982; 57: 243-8
90. Talmadge RJ, Castro MJ, Apple Jr DF, et al. Phenotypic adaptations in human muscle fibers 6 and 24 wk after spinal cord injury. *J Appl Physiol* 2002; 92: 147-54
91. Talmadge RJ, Roy RR, Edgerton VR. Prominence of myosin heavy chain hybrid fibers in soleus muscle of spinal cord-transected rats. *J Appl Physiol* 1995; 78: 1256-65
92. Shields RK. Fatigability, relaxation properties, and electromyographic responses of the human paralyzed soleus muscle. *J Neurophysiol* 1995; 73: 2195-206
93. Shields RK, Law LF, Reiling B, et al. Effects of electrically induced fatigue on the twitch and tetanus of paralyzed soleus muscle in humans. *J Appl Physiol* 1997; 82: 1499-507
94. Rochester L, Barron MJ, Chandler CS, et al. Influence of electrical stimulation of the tibialis anterior muscle in paraplegic subjects, 2: morphological and histochemical properties. *Paraplegia* 1995; 33: 514-22
95. Spungen AM, Bauman WA, Wang J, et al. Measurement of body fat in individuals with tetraplegia: a comparison of eight clinical methods. *Paraplegia* 1995; 33: 402-8
96. McKinley WO, Jackson AB, Cardenas DD, et al. Long-term medical complications after traumatic spinal cord injury: a regional model systems analysis. *Arch Phys Med Rehabil* 1999; 80: 1402-10
97. Claus-Walker J, Carter RE, Compos RJ, et al. Hypercalcemia in early traumatic quadriplegia. *J Chronic Dis* 1975; 28: 81-90
98. Chantraine A, Nussgens B, Lapiere CM. Bone remodeling during the development of osteoporosis in paraplegia. *Calcif Tissue Int* 1986; 38: 323-7
99. Uebelhart D, Demiaux-Domenech B, Roth M, et al. Bone metabolism in spinal cord injured individuals and in others who have prolonged immobilisation: a review. *Paraplegia* 1995; 33: 669-73
100. Ragnarsson KT, Sell GH. Lower extremity fractures after spinal cord injury: a retrospective study. *Arch Phys Med Rehabil* 1981; 62: 418-23
101. Anderson J, Breidahl P. Cartilage atrophy following spinal cord damage. *Australas Radiol* 1981; 25: 98-103
102. Khan MA, Kushner I, Freehafer AA. Sacroiliac joint abnormalities in paraplegics. *Ann Rheum Dis* 1979; 38: 317-9
103. Wright V, Catterall RD, Cook JB. Bone and joint changes in paraplegic men. *Ann Rheum Dis* 1965; 24: 419-31
104. Wylie EJ, Chakera TM. Degenerative joint abnormalities in patients with paraplegia of duration greater than 20 years. *Paraplegia* 1988; 26: 101-6
105. Le CT. Survival from spinal cord injury. *J Chronic Dis* 1982; 35: 487-92
106. Kennedy EJ, Stover SL. Spinal cord injury: the facts and figures. Birmingham (AL): US National Spinal Cord Injury Statistical Center, University of Alabama, 1986
107. Whiteneck G. Learning from empirical investigations. In: Menter R, Whiteneck G, editor. *Perspectives on aging with spinal cord injury*. New York: Demos Publishers, 1992
108. Bauman WA, Spungen AM, Raza M, et al. Coronary artery disease: metabolic risk factors and latent disease in individuals with paraplegia. *Mt Sinai J Med* 1992; 59: 163-8
109. Bauman WA, Raza M, Spungen AM, et al. Cardiac stress testing with thallium-201 imaging reveals silent ischemia in individuals with paraplegia. *Arch Phys Med Rehabil* 1994; 75: 946-50
110. Bauman WA, Raza M, Chayes Z, et al. Tomographic thallium-201 myocardial perfusion imaging after intravenous dipyridamole in asymptomatic subjects with quadriplegia. *Arch Phys Med Rehabil* 1993; 74: 740-4
111. Karlsson AK. Insulin resistance and sympathetic function in high spinal cord injury. *Spinal Cord* 1999; 37: 494-500
112. Karlsson AK, Attvall S, Jansson PA, et al. Influence of the sympathetic nervous system on insulin sensitivity and adipose tissue metabolism: a study in spinal cord-injured subjects. *Metabolism* 1995; 44: 52-8
113. Maki KC, Briones ER, Langbein WE, et al. Associations between serum lipids and indicators of adiposity in men with spinal cord injury. *Paraplegia* 1995; 33: 102-9

114. Zlotolow SP, Levy E, Bauman WA. The serum lipoprotein profile in veterans with paraplegia: the relationship to nutritional factors and body mass index. *J Am Paraplegia Soc* 1992; 15: 158-62
115. Bauman WA, Adkins RH, Spungen AM, et al. Is immobilization associated with an abnormal lipoprotein profile? Observations from a diverse cohort. *Spinal Cord* 1999; 37: 485-93
116. Denke MA, Sempos CT, Grundy SM. Excess body weight: an underrecognized contributor to high blood cholesterol levels in white American men. *Arch Intern Med* 1993; 153: 1093-103
117. Spungen AM, Bauman WA, Wang J, et al. Measurement of body fat in individuals with tetraplegia: a comparison of eight clinical methods. *Paraplegia* 1995; 33: 402-8
118. Washburn RA, Figoni SF. High density lipoprotein cholesterol in individuals with spinal cord injury: the potential role of physical activity. *Spinal Cord* 1999; 37: 685-95
119. Zhong YG, Levy E, Bauman WA. The relationships among serum uric acid, plasma insulin, and serum lipoprotein levels in subjects with spinal cord injury. *Horm Metab Res* 1995; 27: 283-6
120. Grundy SM. Atherogenic dyslipidemia: lipoprotein abnormalities and implications for therapy. *Am J Cardiol* 1995; 75: 45B-52B
121. Aksnes AK, Hjeltne N, Wahlstrom EO, et al. Intact glucose transport in morphologically altered denervated skeletal muscle from quadriplegic patients. *Am J Physiol* 1996; 271: E593-600
122. Bauman WA, Kahn NN, Grimm DR, et al. Risk factors for atherogenesis and cardiovascular autonomic function in persons with spinal cord injury. *Spinal Cord* 1999; 37: 601-16
123. Jones LM, Goulding A, Gerrard DF. DEXA: a practical and accurate tool to demonstrate total and regional bone loss, lean tissue loss and fat mass gain in paraplegia. *Spinal Cord* 1998; 36: 637-40
124. Bauman WA, Adkins RH, Spungen AM, et al. The effect of residual neurological deficit on oral glucose tolerance in persons with chronic spinal cord injury. *Spinal Cord* 1999; 37: 765-71
125. Duckworth WC, Solomon SS, Jallepalli P, et al. Glucose intolerance due to insulin resistance in patients with spinal cord injuries. *Diabetes* 1980; 29: 906-10
126. Kuhne S, Hammon HM, Bruckmaier RM, et al. Growth performance, metabolic and endocrine traits, and absorptive capacity in neonatal calves fed either colostrum or milk replacer at two levels. *J Anim Sci* 2000; 78: 609-20
127. Burstein R, Zeilig G, Royburt M, et al. Insulin resistance in paraplegics: effect of one bout of acute exercise. *Int J Sports Med* 1996; 17: 272-6
128. Cominacini L, Zocca I, Garbin U, et al. High-density lipoprotein composition in obesity: interrelationships with plasma insulin levels and body weight. *Int J Obes* 1988; 12: 343-52
129. Hashimoto R, Adachi H, Tsuruta M, et al. Association of hyperinsulinemia and serum free fatty acids with serum high density lipoprotein-cholesterol. *J Atheroscler Thromb* 1995; 2: 53-9
130. Jeppesen J, Facchini FS, Reaven GM. Individuals with high total cholesterol/HDL cholesterol ratios are insulin resistant. *J Intern Med* 1998; 243: 293-8
131. King ML, Lichtman SW, Pellicone JT, et al. Exertional hypotension in spinal cord injury. *Chest* 1994; 106: 1166-71
132. Kessler KM, Pina I, Green B, et al. Cardiovascular findings in quadriplegic and paraplegic patients and in normal subjects. *Am J Cardiol* 1986; 58: 525-30
133. Cooper IV G, Tomanek RJ. Load regulation of the structure, composition, and function of mammalian myocardium. *Circ Res* 1982; 50: 788-98
134. Nash MS, Bilsker S, Marcillo AE, et al. Reversal of adaptive left ventricular atrophy following electrically-stimulated exercise training in human tetraplegics. *Paraplegia* 1991; 29: 590-9
135. Hopman MT. Circulatory responses during arm exercise in individuals with paraplegia. *Int J Sports Med* 1994; 15: 126-31
136. Hopman MT, van Asten WN, Oeseburg B. Changes in blood flow in the common femoral artery related to inactivity and muscle atrophy in individuals with long-standing paraplegia. *Adv Exp Med Biol* 1996; 388: 379-83
137. Taylor PN, Ewins DJ, Fox B, et al. Limb blood flow, cardiac output and quadriceps muscle bulk following spinal cord injury and the effect of training for the Odstock functional electrical stimulation standing system. *Paraplegia* 1993; 31: 303-10
138. Hjeltne N, Vokac Z. Circulatory strain in everyday life of paraplegics. *Scand J Rehabil Med* 1979; 11: 67-73
139. Boudaoud L, Roussi J, Lortat-Jacob S, et al. Endothelial fibrinolytic reactivity and the risk of deep venous thrombosis after spinal cord injury. *Spinal Cord* 1997; 35: 151-7
140. Brackett NL, Nash MS, Lynne CM. Male fertility following spinal cord injury: facts and fiction. *Phys Ther* 1996; 76: 1221-31
141. Campagnolo DI, Bartlett JA, Keller SE. Influence of neurological level on immune function following spinal cord injury: a review. *J Spinal Cord Med* 2000; 23: 121-8
142. Campagnolo DI, Keller SE, DeLisa JA, et al. Alteration of immune system function in tetraplegics. A pilot study. *Am J Phys Med Rehabil* 1994; 73: 387-93
143. Cruse JM, Keith JC, Bryant Jr ML, et al. Immune system-neuroendocrine dysregulation in spinal cord injury. *Immunol Res* 1996; 15: 306-14
144. Cruse JM, Lewis RE, Bishop GR, et al. Neuroendocrine-immune interactions associated with loss and restoration of immune system function in spinal cord injury and stroke patients. *Immunol Res* 1992; 11: 104-16
145. Cruse JM, Lewis RE, Dilioglou S, et al. Review of immune function, healing of pressure ulcers, and nutritional status in patients with spinal cord injury. *J Spinal Cord Med* 2000; 23: 129-35
146. Cruse JM, Lewis RE, Roe DL, et al. Facilitation of immune function, healing of pressure ulcers, and nutritional status in spinal cord injury patients. *Exp Mol Pathol* 2000; 68: 38-54
147. Cruse JM, Lewis Jr RE, Bishop GR, et al. Decreased immune reactivity and neuroendocrine alterations related to chronic stress in spinal cord injury and stroke patients. *Pathobiology* 1993; 61: 183-92
148. Kliesch WF, Cruse JM, Lewis RE, et al. Restoration of depressed immune function in spinal cord injury patients receiving rehabilitation therapy. *Paraplegia* 1996; 34: 82-90
149. Nash MS. Exercise and immunology. *Med Sci Sports Exerc* 1994; 26: 125-7
150. Nash MS. Known and plausible modulators of depressed immune functions following spinal cord injuries. *J Spinal Cord Med* 2000; 23: 111-20
151. Qian T, Campagnolo D, Kirshblum S. High-dose methylprednisolone may do more harm for spinal cord injury. *Med Hypotheses* 2000; 55: 452-3
152. Figoni SF. Cardiovascular and haemodynamic responses to tilting and to standing in tetraplegic patients: a review. *Paraplegia* 1984; 22: 99-109

153. Minaire P. Immobilization osteoporosis: a review. *Clin Rheumatol* 1989; 8 Suppl. 2: 95-103
154. Gass GC, Camp EM, Nadel ER, et al. Rectal and rectal vs esophageal temperatures in paraplegic men during prolonged exercise. *J Appl Physiol* 1988; 64: 2265-71
155. Hopman MT, Oeseburg B, Binkhorst RA. Cardiovascular responses in paraplegic subjects during arm exercise. *Eur J Appl Physiol* 1992; 65: 73-8
156. Hopman MT, Oeseburg B, Binkhorst RA. The effect of an anti-G suit on cardiovascular responses to exercise in persons with paraplegia. *Med Sci Sports Exerc* 1992; 24: 984-90
157. Menard MR, Hahn G. Acute and chronic hypothermia in a man with spinal cord injury: environmental and pharmacologic causes. *Arch Phys Med Rehabil* 1991; 72: 421-4
158. Bloomfield SA, Jackson RD, Mysiw WJ. Catecholamine response to exercise and training in individuals with spinal cord injury. *Med Sci Sports Exerc* 1994; 26: 1213-9
159. Hooker SP, Greenwood JD, Hatae DT, et al. Oxygen uptake and heart rate relationship in persons with spinal cord injury. *Med Sci Sports Exerc* 1993; 25: 1115-9
160. Irizawa M, Yamasaki M, Muraki S, et al. Relationship between heart rate and oxygen uptake during submaximal arm cranking in paraplegics and quadriplegics. *Ann Physiol Anthropol* 1994; 13: 275-80
161. Schmid A, Huonker M, Stahl F, et al. Free plasma catecholamines in spinal cord injured persons with different injury levels at rest and during exercise. *J Auton Nerv Syst* 1998; 68: 96-100
162. Gass GC, Camp EM. The maximum physiological responses during incremental wheelchair and arm cranking exercise in male paraplegics. *Med Sci Sports Exerc* 1984; 16: 355-9
163. Schneider DA, Sedlock DA, Gass E, et al. VO₂peak and the gas-exchange anaerobic threshold during incremental arm cranking in able-bodied and paraplegic men. *Eur J Appl Physiol* 1999; 80: 292-7
164. Hjeltnes N. Oxygen uptake and cardiac output in graded arm exercise in paraplegics with low level spinal lesions. *Scand J Rehabil Med* 1977; 9: 107-13
165. Hjeltnes N. Cardiorespiratory capacity in tetra- and paraplegia shortly after injury. *Scand J Rehabil Med* 1986; 18: 65-70
166. Hopman MT, Pistorius M, Kamerbeek IC, et al. Cardiac output in paraplegic subjects at high exercise intensities. *Eur J Appl Physiol* 1993; 66: 531-5
167. Van Loan MD, McCluer S, Loftin JM, et al. Comparison of physiological responses to maximal arm exercise among able-bodied, paraplegics and quadriplegics. *Paraplegia* 1987; 25: 397-405
168. Houtman S, Thielen JJ, Binkhorst RA, et al. Effect of a pulsating anti-gravity suit on peak exercise performance in individual with spinal cord injuries. *Eur J Appl Physiol* 1999; 79: 202-4
169. Schmid A, Huonker M, Barturen JM, et al. Catecholamines, heart rate, and oxygen uptake during exercise in persons with spinal cord injury. *J Appl Physiol* 1998; 85: 635-41
170. Davis GM, Shephard RJ. Cardiorespiratory fitness in highly active versus inactive paraplegics. *Med Sci Sports Exerc* 1988; 20: 463-8
171. Janssen TW, van Oers CA, van der Woude LH, et al. Physical strain in daily life of wheelchair users with spinal cord injuries. *Med Sci Sports Exerc* 1994; 26: 661-70
172. Franklin BA. Exercise testing, training and arm ergometry. *Sports Med* 1985; 2: 100-19
173. Jacobs PL, Nash MS, Klose KJ, et al. Evaluation of a training program for persons with SCI paraplegia using the Parastep I ambulation system, part 2: effects on physiological responses to peak arm ergometry. *Arch Phys Med Rehabil* 1997; 78: 794-8
174. Jacobs PL, Nash MS, Rusinowski JW. Circuit training provides cardiorespiratory and strength benefits in persons with paraplegia. *Med Sci Sports Exerc* 2001; 33: 711-7
175. Nash MS, Jacobs PL, Mendez AJ, et al. Circuit resistance training improves the atherogenic lipid profiles of persons with chronic paraplegia. *J Spinal Cord Med* 2001; 24: 2-9
176. Davis GM, Kofsky PR, Kelsey JC, et al. Cardiorespiratory fitness and muscular strength of wheelchair users. *CMAJ* 1981; 125: 1317-23
177. Davis GM, Shephard RJ, Leenen FH. Cardiac effects of short term arm crank training in paraplegics: echocardiographic evidence. *Eur J Appl Physiol* 1987; 56: 90-6
178. DiCarlo SE, Supp MD, Taylor HC. Effect of arm ergometry training on physical work capacity of individuals with spinal cord injuries. *Phys Ther* 1983; 63: 1104-7
179. Gass GC, Watson J, Camp EM, et al. The effects of physical training on high level spinal lesion patients. *Scand J Rehabil Med* 1980; 12: 61-5
180. Hooker SP, Wells CL. Effects of low- and moderate-intensity training in spinal cord-injured persons. *Med Sci Sports Exerc* 1989; 21: 18-22
181. Knutsson E, Lewenhaupt-Olsson E, Thorsen M. Physical work capacity and physical conditioning in paraplegic patients. *Paraplegia* 1973; 11: 205-16
182. Miles DS, Sawka MN, Wilde SW, et al. Pulmonary function changes in wheelchair athletes subsequent to exercise training. *Ergonomics* 1982; 25: 239-46
183. Pachalski A, Mekarski T. Effect of swimming on increasing of cardio-respiratory capacity in paraplegics. *Paraplegia* 1980; 18: 190-6
184. Pollock ML, Miller HS, Linnerud AC, et al. Arm pedaling as an endurance training regimen for the disabled. *Arch Phys Med Rehabil* 1974; 55: 418-24
185. Sedlock DA, Knowlton RG, Fitzgerald PI. The effects of arm crank training on the physiological responses to submaximal wheelchair ergometry. *Eur J Appl Physiol* 1988; 57: 55-9
186. Taylor AW, McDonnell E, Brassard L. The effects of an arm ergometer training programme on wheelchair subjects. *Paraplegia* 1986; 24: 105-14
187. Yim SY, Cho KJ, Park CI, et al. Effect of wheelchair ergometer training on spinal cord-injured paraplegics. *Yonsei Med J* 1993; 34: 278-86
188. DiCarlo SE. Effect of arm ergometry training on wheelchair propulsion endurance of individuals with quadriplegia. *Phys Ther* 1988; 68: 40-4
189. Drory Y, Ohry A, Brooks ME, et al. Arm crank ergometry in chronic spinal cord injured patients. *Arch Phys Med Rehabil* 1990; 71: 389-92
190. Pollock ML, Wilmore JH. Exercise in health and disease. Philadelphia (PA): WB Saunders, 1990: 670-1
191. Wicks JR, Oldridge NB, Cameron BJ, et al. Arm cranking and wheelchair ergometry in elite spinal cord-injured athletes. *Med Sci Sports Exerc* 1983; 15: 224-31
192. Fignon SF. Spinal cord injury. In: Durstine JL, editor. Exercise management for persons with chronic diseases and disabilities. Champaign (IL): Human Kinetics, 1997: 175-9
193. Lockette KF, Keyes AM. Conditioning with physical disabilities. Champaign (IL): Human Kinetics, 1994
194. Miller PD. Fitness programming and physical disability. Champaign (IL): Human Kinetics, 1995

195. Escobedo EM, Hunter JC, Hollister MC, et al. MR imaging of rotator cuff tears in individuals with paraplegia. *AJR Am J Roentgenol* 1997; 168: 919-23
196. Gellman H, Sie I, Waters RL. Late complications of the weight-bearing upper extremity in the paraplegic patient. *Clin Orthop* 1988; 233: 132-5
197. Goldstein B, Young J, Escobedo EM. Rotator cuff repairs in individuals with paraplegia. *Am J Phys Med Rehabil* 1997; 76: 316-22
198. Nichols PJ, Norman PA, Ennis JR. Wheelchair user's shoulder? Shoulder pain in patients with spinal cord lesions. *Scand J Rehabil Med* 1979; 11: 29-32
199. Pentland WE, Twomey LT. The weight-bearing upper extremity in women with long term paraplegia. *Paraplegia* 1991; 29: 521-30
200. Pentland WE, Twomey LT. Upper limb function in persons with long term paraplegia and implications for independence: part I. *Paraplegia* 1994; 32: 211-8
201. Burnham RS, May L, Nelson E, et al. Shoulder pain in wheelchair athletes: the role of muscle imbalance. *Am J Sports Med* 1993; 21: 238-42
202. Curtis KA, Roach KE, Applegate EB, et al. Reliability and validity of the Wheelchair User's Shoulder Pain Index (WUS-PI). *Paraplegia* 1995; 33: 595-601
203. Nilsson S, Staff PH, Pruett ED. Physical work capacity and the effect of training on subjects with long-standing paraplegia. *Scand J Rehabil Med* 1975; 7: 51-6
204. Davis GM, Shephard RJ. Strength training for wheelchair users. *Br J Sports Med* 1990; 24: 25-30
205. Cooney MM, Walker JB. Hydraulic resistance exercise benefits cardiovascular fitness of spinal cord injured. *Med Sci Sports Exerc* 1986; 18: 522-5
206. Olenik LM, Laskin JJ, Burnham R, et al. Efficacy of rowing, backward wheeling and isolated scapular retractor exercise as remedial strength activities for wheelchair users: application of electromyography. *Paraplegia* 1995; 33: 148-52
207. Gettman LR, Ayres JJ, Pollock ML, et al. The effect of circuit weight training on strength, cardiorespiratory function, and body composition of adult men. *Med Sci Sports* 1978; 10: 171-6
208. Nash MS, Jacobs PL, Woods JM, et al. A comparison of 2 circuit exercise training techniques for eliciting matched metabolic responses in persons with paraplegia. *Arch Phys Med Rehabil* 2002; 83: 201-9
209. Dudley GA, Castro MJ, Rogers S, et al. A simple means of increasing muscle size after spinal cord injury: a pilot study. *Eur J Appl Physiol Occup Physiol* 1999; 80: 394-6
210. Ficoni SF, Glaser RM, Rodgers MM, et al. Acute hemodynamic responses of spinal cord injured individuals to functional neuromuscular stimulation-induced knee extension exercise. *J Rehabil Res Dev* 1991; 28: 9-18
211. Gruner JA, Glaser RM, Feinberg SD, et al. A system for evaluation and exercise-conditioning of paralyzed leg muscles. *J Rehabil R D* 1983; 20: 21-30
212. Rodgers MM, Glaser RM, Ficoni SF, et al. Musculoskeletal responses of spinal cord injured individuals to functional neuromuscular stimulation-induced knee extension exercise training. *J Rehabil Res Dev* 1991; 28: 19-26
213. Billian C, Gorman PH. Upper extremity applications of functional neuromuscular stimulation. *Assist Technol* 1992; 4: 31-9
214. Bryden AM, Memberg WD, Crago PE. Electrically stimulated elbow extension in persons with C5/C6 tetraplegia: a functional and physiological evaluation. *Arch Phys Med Rehabil* 2000; 81: 80-8
215. Klefbeck B, Mattsson E, Weinberg J. The effect of trunk support on performance during arm ergometry in patients with cervical cord injuries. *Paraplegia* 1996; 34: 167-72
216. Li GL, Farooque M, Holtz A, et al. Expression of the ubiquitin carboxyl-terminal hydrolase PGP 9.5 in axons following spinal cord compression trauma. An immunohistochemical study in the rat. *APMIS* 1997; 105: 384-90
217. Mulcahey MJ, Betz RR, Smith BT, et al. Implanted functional electrical stimulation hand system in adolescents with spinal injuries: an evaluation. *Arch Phys Med Rehabil* 1997; 78: 597-607
218. Scremin AM, Kurta L, Gentili A, et al. Increasing muscle mass in spinal cord injured persons with a functional electrical stimulation exercise program. *Arch Phys Med Rehabil* 1999; 80: 1531-6
219. Glaser RM. Physiologic aspects of spinal cord injury and functional neuromuscular stimulation. *Cent Nerv Syst Trauma* 1986; 3: 49-62
220. Hooker SP, Ficoni SF, Glaser RM, et al. Physiologic responses to prolonged electrically stimulated leg-cycle exercise in the spinal cord injured. *Arch Phys Med Rehabil* 1990; 71: 863-9
221. Ragnarsson KT, Pollack S, O'Daniel Jr W, et al. Clinical evaluation of computerized functional electrical stimulation after spinal cord injury: a multicenter pilot study. *Arch Phys Med Rehabil* 1988; 69: 672-7
222. Raymond J, Davis GM, Climstein M, et al. Cardiorespiratory responses to arm cranking and electrical stimulation leg cycling in people with paraplegia. *Med Sci Sports Exerc* 1999; 31: 822-8
223. Krauss JC, Robergs RA, Depaape JL, et al. Effects of electrical stimulation and upper body training after spinal cord injury. *Med Sci Sports Exerc* 1993; 25: 1054-61
224. Mutton DL, Scremin AM, Barstow TJ, et al. Physiologic responses during functional electrical stimulation leg cycling and hybrid exercise in spinal cord injured subjects. *Arch Phys Med Rehabil* 1997; 78: 712-8
225. Phillips W, Burkett LN. Arm crank exercise with static leg FNS in persons with spinal cord injury. *Med Sci Sports Exerc* 1995; 27: 530-5
226. Cameron T, Broton JG, Needham-Shropshire B, et al. An upper body exercise system incorporating resistive exercise and neuromuscular electrical stimulation (NMS). *J Spinal Cord Med* 1998; 21: 1-6
227. Needham-Shropshire BM, Broton JG, Cameron TL, et al. Improved motor function in tetraplegics following neuromuscular stimulation-assisted arm ergometry. *J Spinal Cord Med* 1997; 20: 49-55
228. Phillips CA. Functional electrical stimulation and lower extremity bracing for ambulation exercise of the spinal cord injured individual: a medically prescribed system. *Phys Ther* 1989; 69: 842-9
229. Phillips CA, Gallimore JJ, Hendershot DM. Walking when utilizing a sensory feedback system and an electrical muscle stimulation gait orthosis. *Med Eng Phys* 1995; 17: 507-13
230. Brissot R, Gallien P, Le Bot MP, et al. Clinical experience with functional electrical stimulation-assisted gait with Parastep in spinal cord-injured patients. *Spine* 2000; 25: 501-8
231. Guest RS, Klose KJ, Needham-Shropshire BM, et al. Evaluation of a training program for persons with SCI paraplegia using the Parastep 1 ambulation system, part 4: effect on physical self-

- concept and depression. *Arch Phys Med Rehabil* 1997; 78: 804-7
232. Needham-Shropshire BM, Broton JG, Klose KJ, et al. Evaluation of a training program for persons with SCI paraplegia using the Parastep 1 ambulation system, part 3: lack of effect on bone mineral density. *Arch Phys Med Rehabil* 1997; 78: 799-803
 233. Phillips CA. Medical criteria for active physical therapy: physician guidelines for patient participation in a program of functional electrical rehabilitation. *Am J Phys Med* 1987; 66: 269-86
 234. Dimitrijevic MM. Mesh-glove: 1. A method for whole-hand electrical stimulation in upper motor neuron dysfunction. *Scand J Rehabil Med* 1994; 26: 183-6
 235. Kameyama J, Handa Y, Hoshimiya N, et al. Restoration of shoulder movement in quadriplegic and hemiplegic patients by functional electrical stimulation using percutaneous multiple electrodes. *Tohoku J Exp Med* 1999; 187: 329-37
 236. Mulcahey MJ, Smith BT, Betz RR. Evaluation of the lower motor neuron integrity of upper extremity muscles in high level spinal cord injury. *Spinal Cord* 1999; 37: 585-91
 237. Peckham PH, Creasey GH. Neural prostheses: clinical applications of functional electrical stimulation in spinal cord injury. *Paraplegia* 1992; 30: 96-101
 238. Scott TR, Peckham PH, Keith MW. Upper extremity neuroprostheses using functional electrical stimulation. *Baillieres Clin Neurol* 1995; 4: 57-75
 239. Graupe D. An overview of the state of the art of noninvasive FES for independent ambulation by thoracic level paraplegics. *Neurol Res* 2002; 24: 431-42
 240. Ragnarsson KT. Physiologic effects of functional electrical stimulation-induced exercises in spinal cord-injured individuals. *Clin Orthop* 1988; 233: 53-63
 241. Petrofsky JS, Phillips CA. The use of functional electrical stimulation for rehabilitation of spinal cord injured patients. *Cent Nerv Syst Trauma* 1984; 1: 57-74
 242. Hooker SP, Figoni SF, Rodgers MM, et al. Physiologic effects of electrical stimulation leg cycle exercise training in spinal cord injured persons. *Arch Phys Med Rehabil* 1992; 73: 470-6
 243. Hooker SP, Scremin AM, Mutton DL, et al. Peak and submaximal physiologic responses following electrical stimulation leg cycle ergometer training. *J Rehabil Res Dev* 1995; 32: 361-6
 244. Barstow TJ, Scremin AM, Mutton DL, et al. Gas exchange kinetics during functional electrical stimulation in subjects with spinal cord injury. *Med Sci Sports Exerc* 1995; 27: 1284-91
 245. Barstow TJ, Scremin AM, Mutton DL, et al. Changes in gas exchange kinetics with training in patients with spinal cord injury. *Med Sci Sports Exerc* 1996; 28: 1221-8
 246. Gerrits HL, de Haan A, Sargeant AJ, et al. Peripheral vascular changes after electrically stimulated cycle training in people with spinal cord injury. *Arch Phys Med Rehabil* 2001; 82: 832-9
 247. Schmidt-Trucksass A, Schmid A, Brunner C, et al. Arterial properties of the carotid and femoral artery in endurance-trained and paraplegic subjects. *J Appl Physiol* 2000; 89: 1956-63
 248. BeDell KK, Scremin AM, Perell KL, et al. Effects of functional electrical stimulation-induced lower extremity cycling on bone density of spinal cord-injured patients. *Am J Phys Med Rehabil* 1996; 75: 29-34
 249. Mohr T, Podenphant J, Biering-Sorensen F, et al. Increased bone mineral density after prolonged electrically induced cycle training of paralyzed limbs in spinal cord injured man. *Calcif Tissue Int* 1997; 61: 22-5
 250. Leeds EM, Klose KJ, Ganz W, et al. Bone mineral density after bicycle ergometry training. *Arch Phys Med Rehabil* 1990; 71: 207-9
 251. Hjeltnes N, Aksnes AK, Birkeland KI, et al. Improved body composition after 8 wk of electrically stimulated leg cycling in tetraplegic patients. *Am J Physiol* 1997; 273: R1072-9
 252. Hjeltnes N, Galuska D, Bjornholm M, et al. Exercise-induced overexpression of key regulatory proteins involved in glucose uptake and metabolism in tetraplegic persons: molecular mechanism for improved glucose homeostasis. *FASEB J* 1998; 12: 1701-12
 253. Mohr T, Dela F, Handberg A, et al. Insulin action and long-term electrically induced training in individuals with spinal cord injuries. *Med Sci Sports Exerc* 2001; 33: 1247-52
 254. Gallien P, Brisot R, Eyssette M, et al. Restoration of gait by functional electrical stimulation for spinal cord injured patients. *Paraplegia* 1995; 33: 660-4
 255. Bajd T, Andrews BJ, Kralj A, et al. Restoration of walking in patients with incomplete spinal cord injuries by use of surface electrical stimulation: preliminary results. *Prosthet Orthot Int* 1985; 9: 109-11
 256. Bajd T, Kralj A, Stefancic M, et al. Use of functional electrical stimulation in the lower extremities of incomplete spinal cord injured patients. *Artif Organs* 1999; 23: 403-9
 257. Bajd T, Munih M, Savrin R, et al. Dermatome electrical stimulation as a therapeutic ambulatory aid for incomplete spinal cord injured patients. *Artif Organs* 2002; 26: 260-2
 258. Barbeau H, Ladouceur M, Mirbagheri MM, et al. The effect of locomotor training combined with functional electrical stimulation in chronic spinal cord injured subjects: walking and reflex studies. *Brain Res Brain Res Rev* 2002; 40: 274-91
 259. Field-Fote E. Spinal cord stimulation facilitates functional walking in a chronic, incomplete spinal cord injured subject. *Spinal Cord* 2002; 40: 428
 260. Granat M, Keating JF, Smith AC, et al. The use of functional electrical stimulation to assist gait in patients with incomplete spinal cord injury. *Disabil Rehabil* 1992; 14: 93-7
 261. Ladouceur M, Barbeau H. Functional electrical stimulation-assisted walking for persons with incomplete spinal injuries: longitudinal changes in maximal overground walking speed. *Scand J Rehabil Med* 2000; 32: 28-36
 262. Ladouceur M, Barbeau H. Functional electrical stimulation-assisted walking for persons with incomplete spinal injuries: changes in the kinematics and physiological cost of overground walking. *Scand J Rehabil Med* 2000; 32: 72-9
 263. Stein RB, Belanger M, Wheeler G, et al. Electrical systems for improving locomotion after incomplete spinal cord injury: an assessment. *Arch Phys Med Rehabil* 1993; 74: 954-9
 264. Graupe D, Kohn KH. Transcutaneous functional neuromuscular stimulation of certain traumatic complete thoracic paraplegics for independent short-distance ambulation. *Neurol Res* 1997; 19: 323-33
 265. Jaeger RJ. Lower extremity applications of functional neuromuscular stimulation. *Assist Technol* 1992; 4: 19-30
 266. Erickson RP. Autonomic hyperreflexia: pathophysiology and medical management. *Arch Phys Med Rehabil* 1980; 61: 431-40
 267. Comarr AE, Eltorai I. Autonomic dysreflexia/hyperreflexia. *J Spinal Cord Med* 1997; 20: 345-54

268. Chancellor MB, Erhard MJ, Hirsch IH, et al. Prospective evaluation of terazosin for the treatment of autonomic dysreflexia. *J Urol* 1994; 151: 111-3
269. Steinberger RE, Ohl DA, Bennett CJ, et al. Nifedipine pretreatment for autonomic dysreflexia during electroejaculation. *Urology* 1990; 36: 228-31
270. Vaidyanathan S, Soni BM, Sett P, et al. Pathophysiology of autonomic dysreflexia: long-term treatment with terazosin in adult and paediatric spinal cord injury patients manifesting recurrent dysreflexic episodes. *Spinal Cord* 1998; 36: 761-70
271. Webborn AD. 'Boosting' performance in disability sport. *Br J Sports Med* 1999; 33: 74-5
272. Curtis KA, Drysdale GA, Lanza RD, et al. Shoulder pain in wheelchair users with tetraplegia and paraplegia. *Arch Phys Med Rehabil* 1999; 80: 453-7
273. Ballinger DA, Rintala DH, Hart KA. The relation of shoulder pain and range-of-motion problems to functional limitations, disability, and perceived health of men with spinal cord injury: a multifaceted longitudinal study. *Arch Phys Med Rehabil* 2000; 81: 1575-81
274. Bayley JC, Cochran TP, Sledge CB. The weight-bearing shoulder: the impingement syndrome in paraplegics. *J Bone Joint Surg Am* 1987; 69: 676-8
275. King ML, Freeman DM, Pellicone JT, et al. Exertional hypotension in thoracic spinal cord injury: case report. *Paraplegia* 1992; 30: 261-6
276. Lopes P, Figoni SF, Perakash I. Upper limb exercise effect on tilt tolerance during orthostatic training of patients with spinal cord injury. *Arch Phys Med Rehabil* 1984; 65: 251-3
277. Gerner HJ, Engel P, Gass GC, et al. The effects of sauna on tetraplegic and paraplegic subjects. *Paraplegia* 1992; 30: 410-9
278. Ishii K, Yamasaki M, Muraki S, et al. Effects of upper limb exercise on thermoregulatory responses in patients with spinal cord injury. *Appl Human Sci* 1995; 14: 149-54
279. Sawka MN, Latzka WA, Pandolf KB. Temperature regulation during upper body exercise: able-bodied and spinal cord injured. *Med Sci Sports Exerc* 1989; 21: S132-40
280. Muraki S, Yamasaki M, Ishii K, et al. Effect of arm cranking exercise on skin blood flow of lower limb in people with injuries to the spinal cord. *Eur J Appl Physiol Occup Physiol* 1995; 71: 28-32
281. Muraki S, Yamasaki M, Ishii K, et al. Relationship between core temperature and skin blood flux in lower limbs during prolonged arm exercise in persons with spinal cord injury. *Eur J Appl Physiol* 1996; 72: 330-4
282. Price MJ, Campbell IG. Thermoregulatory responses of spinal cord injured and able-bodied athletes to prolonged upper body exercise and recovery. *Spinal Cord* 1999; 37: 772-9

Correspondence and offprints: Dr *Patrick L. Jacobs*, The Miami Project to Cure Paralysis, University of Miami School of Medicine, 1095 Northwest 14th Terrace, Miami, R-48, FL 33136, USA.
E-mail: pjacobs@miami.edu