

# the BRIEFING

Exeter HS&DR Evidence Synthesis Centre November 2023

## Remote monitoring for long-term physical health conditions: An Evidence and Gap Map

**M**onitoring a patient remotely allows healthcare professionals to understand what is happening to their health between visits and respond to changes in a patient's condition<sup>1</sup>. As an innovative use of technology, remote monitoring could help meet changing demand for health services resulting from an aging population and increasing rates of noncommunicable diseases<sup>1,2</sup>. It could also contribute to the development of a more sustainable healthcare system.

**This is a summary of a project which produced an Evidence and Gap Map (EGM) on the effectiveness, acceptability, and implementation of remote monitoring for long-term physical health conditions. EGMs draw together all the available evidence on a topic, highlighting for which populations or intervention types there is either a lot, or lack, of research. By mapping evidence for different types of remote monitoring, this EGM is intended to support the design and delivery of interventions, and to help prioritise future research.**


It found that:

- ◆ Evidence is focused on remote monitoring of people with a few specific conditions such as heart disease and diabetes, with a lack of research on less common conditions e.g. epilepsy;
- ◆ There are reviews on the effectiveness of remote monitoring for physical and mental health, and health service use, but few which look at the views of carers and healthcare professionals on its acceptability and implementation; and
- ◆ Reviews need to include more detailed descriptions of included remote monitoring interventions.

Remote monitoring:  
*the monitoring of a patient (including self-monitoring), allowing healthcare professionals to assess and manage a patient's condition remotely - without the need for the patient to be seen face-to-face*

### Exeter HS&DR Evidence Synthesis Centre

We are one of three research groups in the UK commissioned by the National Institute of Health Research HS&DR (Health Services & Delivery Research Programme) to conduct syntheses of evidence about the organisation and delivery of healthcare (Project number 16/47/22). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.



FUNDED BY

**NIHR** | National Institute  
for Health Research

## Why did we do this review?

The use of remote monitoring has been driven by the need to deliver health services efficiently. It could allow staff to use their time more effectively e.g. by reducing time spent travelling<sup>3</sup>, and lead to fewer face-to-face appointments, therefore reducing costs<sup>4</sup>.

Remote monitoring also offers many potential benefits to patients. Detecting changes in a patient's condition early means interventions can

be made e.g. changes to medication<sup>5</sup>. Monitoring gives patients an opportunity to learn about their health condition and become more engaged in making decisions about their healthcare<sup>6</sup>.

However, interventions vary widely, including:

- ◆ The technology used to measure health e.g. whether this automatically uploads data;
- ◆ The frequency of contact with healthcare professionals; and
- ◆ The type and content of feedback.

This variation creates challenges for those who are designing or delivering remote monitoring interventions, including policymakers and practitioners. Being able to find and understand evidence relevant to the type of remote monitoring which is being planned or delivered should aid decision-making.

Our aim was to map all the recent, high quality evidence on the use of remote monitoring for long-term physical health conditions, looking at both its effectiveness, and the acceptability and implementation of interventions.




## How did we make this EGM?

**Finding the literature:** We searched ten bibliographic databases for systematic reviews. We also searched the citations and reference lists of included reviews and registries of review protocols.

**Eligibility criteria:** Systematic reviews were included in the map. Included reviews:

- Focused on adult populations with a long-term physical health condition(s).
- Focused on any type of remote monitoring intervention, as long as data was passed to a healthcare professional.
- Contained studies conducted in high income countries.
- Were published in English from 2018 onwards.



**Study selection, data extraction and assessment of study quality:** Studies were screened independently by two reviewers for inclusion. Data extraction and assessment of study quality were then carried out by one reviewer and checked by another. Disagreements were resolved through discussion. We used a standard tool, AMSTAR 2, to assess study quality.

Data were entered into EPPI-Reviewer 4 software and used to construct a web-based interactive map.

## What did we find?

Of the 72 reviews investigating remote monitoring in adults with a long-term physical condition included in the map, 61 focus on effectiveness and 24 on how to implement remote monitoring, including whether it is acceptable to patients, carers and healthcare professionals.

Most reviews were assessed to be of low or critically low quality. Most reviews were rated of lower quality as they did not have a pre-registered protocol.

There were several areas of focus:

- ◆ Most reviews included studies from North America and Europe, with 38 including studies from the UK;
- ◆ The most common health conditions were heart disease, diabetes, and lung conditions;
- ◆ Nurses were the healthcare professional most often involved in interventions;
- ◆ Most feedback was by telephone and contained motivational/educational elements; and
- ◆ Reviews focused on whether remote monitoring affected physical and mental health, and health service use.

There were also gaps in the evidence:

- ◆ There was no or little evidence for some health conditions e.g. epilepsy; and
- ◆ Few reviews contained studies on the acceptability and implementation of remote monitoring for carers and healthcare professionals.

There was a lack of detail in the descriptions of the remote monitoring interventions in the reviews e.g. the type of healthcare professional involved was not reported consistently, or the content of feedback. There was also a lack of consistent reporting on factors (e.g. age, gender, and digital literacy) that could affect effectiveness of remote monitoring interventions.

## The interactive EGM can be accessed [HERE](#) online.

The picture below shows the map which has a typical EGM format: the rows give details of the remote monitoring intervention and the columns show what outcomes were measured by the study.

Each cell shows the studies giving evidence on that particular combination of intervention and outcome, with the different colours of the squares in the cell indicating the quality of the studies. For each study, we have provided an abstract or summary and a link to the original source.

Filters can also be applied to the map, meaning the map only displays evidence for the selected filter. These filters include population categories and type of review.

This is an evidence and gap map of systematic reviews. The rows show intervention categories and the columns indicate outcome categories. Both can be maximised or minimised by clicking the small arrows in the middle row. The cells in the table contain tiles, the number of these indicates the number of reviews in the cell. Different colours represent the study quality; if you hover over the map, you can see the meaning of each colour (key also available in left hand bottom corner of page). The full report, detailed [map instructions](#), and a [glossary](#) are available.

		Outcomes									
		Physical health									Mental health/wellbeing
		Mortality	Blood glucose/ glycaemic control	Blood pressure	Other cardiovascular metrics e.g. oxygen consumption	Detection rate e.g. exacerbation, atrial arrhythmia	Risk of adverse events e.g. stroke	Weight/BMI/waist circumference	General health	Other	Anxiety/ depression
Intervention (characteristics of remote monitoring)	What is monitored	[Grid of colored tiles representing studies and their quality]									
	How is it monitored	[Grid of colored tiles representing studies and their quality]									

## What are the implications of this EGM?

Systematic reviews of the effectiveness, acceptability and implementation of remote monitoring for adults with long-term physical health conditions could provide evidence to support the provision of interventions.

There were several areas of evidence which could be particularly useful in informing the design and delivery of interventions:

- ◆ The effectiveness of remote monitoring for common health conditions such as heart disease, diabetes, and lung disease e.g. COPD;
- ◆ For measuring aspects of health status related to these conditions e.g. blood pressure, heart rate; and
- ◆ Using common devices such as heart rate monitors or blood pressure monitors.

There is a lack of research in some areas where further evidence could support the provision of remote monitoring, including:

- ◆ Evidence on other conditions such as dementia or epilepsy;

- ◆ The views of carers and healthcare professionals on factors which would improve the acceptability of monitoring and aid its implementation.

These gaps indicate a need for more studies. In some cases, this will be primary research, such as randomised controlled trials, to evaluate existing remote monitoring interventions. Where evidence exists, systematic reviews are needed to synthesise this research.

Studies should describe the details of remote monitoring interventions more clearly. A lack of clear reporting made it difficult to tell how similar interventions were in different reviews. Some information on participants e.g. age, digital literacy, was poorly reported yet has implications for effectiveness and health equity.

For this EGM, we had to exclude studies of interventions with multiple components (e.g. education and monitoring), so may have missed some relevant evidence. Whilst more than half of included reviews have serious methodological issues, they may still contain high quality studies.



### Contact Us

Sian de Bell  
s.c.de-bell@exeter.ac.uk

Zhivko Zhelev  
z.zhelev@exeter.ac.uk

Visit our [webpage](#).

@ExEvidSC

Please use the QR code to view the project report online:



## References

- 1 Kvedar J, Coye MJ, Everett W. Connected health: A review of technologies and strategies to improve patient care with telemedicine and telehealth. *Health Affairs* 2014;33:194-9. <https://doi.org/10.1377/hlthaff.2013.0992>
- 2 Government Office for Science, Foresight. *Future of an ageing population*. London: Government Office for Science; 2016
- 3 Davis MM, Freeman M, Kaye J, Vuckovic N, Buckley DI. A systematic review of clinician and staff views on the acceptability of incorporating remote monitoring technology into primary care. *Telemed e-Health* 2014;20:428-38. <https://doi.org/https://dx.doi.org/10.1089/tmj.2013.0166>
- 4 McBain H, Shipley M, Newman S. The impact of self-monitoring in chronic illness on healthcare utilisation: a systematic review of reviews. *BMC Health Serv Res* 2015;15:565. <https://doi.org/10.1186/s12913-015-1221-5>
- 5 Iqbal FM, Lam K, Joshi M, Khan S, Ashrafian H, Darzi A. Clinical outcomes of digital sensor alerting systems in remote monitoring: a systematic review and meta-analysis. *NPJ Digit Med* 2021;4:7. <https://doi.org/10.1038/s41746-020-00378-0>
- 6 Lancaster K, Abuzour A, Khaira M, Mathers A, Chan A, Bui V, et al. The use and effects of Electronic Health Tools for patient self-monitoring and reporting of outcomes following medication use: Systematic review. *J Med Internet Res* 2018;20:e294. <https://doi.org/https://dx.doi.org/10.2196/jmir.9284>
- 7 Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomized studies of healthcare interventions, or both. *BMJ* 2017;358:j4008. <https://doi.org/10.1136/bmj.j4008>